Counsyl

RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Dr. Jeffrey Olliffe
4915 25th Ave NE, Suite 204W
Seattle, WA 98105
Phone: (206) 588-1484
Fax: (206) 466-4696
NPI: 1306838271
Report Date: 09/10/2018

MALE
DONOR 10289
DOB: [redacted]
Ethnicity: Southeast Asian
Sample Type: EDTA Blood
Date of Collection: 08/31/2018
Date Received: 09/04/2018
Date Tested: 09/10/2018
Barcode: 11004212488944
Accession ID: CSL6D9P9KKQJK3N
Indication: Egg or sperm donor

FEMALE
N/A

Foresight™ Carrier Screen

POSITIVE: CARRIER

ABOUT THIS TEST

The **Counsyl Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

RESULTS SUMMARY

Risk Details	DONOR 10289	Partner
Panel Information	Foresight Carrier Screen Universal Panel **(175 conditions tested)**	N/A
POSITIVE: CARRIER Beta-sarcoglycanopathy **Reproductive Risk: 1 in 2,000** **Inheritance: Autosomal Recessive**	**CARRIER*** NM_000232.4(SGCB):c.1A>G(M1?) heterozygote †	The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group. Carrier testing should be considered. See "Next Steps".

†Likely to have a negative impact on gene function.
*Carriers generally do not experience symptoms.

No disease-causing mutations were detected in any other gene tested. A complete list of all conditions tested can be found on page 6.

CLINICAL NOTES

- None

NEXT STEPS

- Carrier testing should be considered for the diseases specified above for the patient's partner, as both parents must be carriers before a child is at high risk of developing the disease.
- Genetic counseling is recommended and patients may wish to discuss any positive results with blood relatives, as there is an increased chance that they are also carriers.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

POSITIVE: CARRIER

Beta-sarcoglycanopathy

Reproductive risk: 1 in 2,000
Risk before testing: < 1 in 1,000,000

Gene: SGCB | **Inheritance Pattern:** Autosomal Recessive

Patient	DONOR 10289	No partner tested
Result	Carrier	N/A
Variant(s)	NM_000232.4(SGCB):c.1A>G(M1?) heterozygote †	N/A
Methodology	Sequencing with copy number analysis	N/A
Interpretation	This individual is a carrier of beta-sarcoglycanopathy. Carriers generally do not experience symptoms.	N/A
Detection rate	>99%	N/A
Exons tested	NM_000232:1-6.	N/A

†Likely to have a negative impact on gene function.

What is Beta-Sarcoglycanopathy?

Beta-sarcoglycanopathy, also known as limb-girdle muscular dystrophy type 2E (LGMD2E), typically causes muscle weakness as a result of a deficiency of the protein, beta-sarcoglycan, in the dystrophin-glycoprotein complex, a component of the muscle system. Symptoms of the disease vary greatly from person to person, even among people in the same family. Some people with the disease can have a mild course where they are nearly asymptomatic, while others may have severe symptoms that can be fatal.

People with beta-sarcoglycanopathy develop symptoms at variable ages, though symptoms tend to first present in childhood. Beta-sarcoglycanopathy does not affect intelligence or mental function; the most common symptom is progressive muscle weakness of the hip, shoulder, and abdomen (proximal muscles). The rate at which the muscles weaken can vary, but many experience progressive weakness to a point where a wheelchair becomes necessary. Other possible features include enlarged calf muscles (calf hypertrophy), contractures, scapular winging (shoulder blade is prominent), and scoliosis. Respiratory complications (~10-30% of individuals) or heart complications (~60-70% of individuals) are also associated with the sarcoglycanopathies, and may be a cause of death.

How common is Beta-Sarcoglycanopathy?

There are numerous types of limb-girdle muscular dystrophy. The estimated prevalence of all types of limb-girdle muscular dystrophy is 1 in 15,000 individuals. Beta-sarcoglycanopathy is rare and its exact incidence is unknown.

How is Beta-Sarcoglycanopathy treated?

There is no cure for beta-sarcoglycanopathy and few effective treatments. Physical therapy is often recommended to retain muscle strength and mobility for as long as possible. Stretching, mechanical aids, or surgery may aid in that goal. As muscles deteriorate, a ventilator may be required to aid breathing. Cardiac surveillance is recommended, and those who develop heart problems should consult with a cardiologist for symptomatic treatments.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

What is the prognosis for a person with Beta-Sarcoglycanopathy?

The outlook for a person with beta-sarcoglycanopathy varies. Generally speaking, the earlier symptoms begin, the faster they progress. However, because symptoms and onset can be variable, prognosis can be variable. People with more severe symptoms can become wheelchair bound in their early teens and die in early adulthood with death usually being due to respiratory and/or cardiac complications.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Methods and Limitations

DONOR 10289 [Foresight Carrier Screen]: Sequencing with copy number analysis, spinal muscular atrophy, and analysis of homologous regions.

Sequencing with copy number analysis

High-throughput sequencing and read depth-based copy number analysis are used to analyze the listed exons, as well as selected intergenic and intronic regions, of the genes in the Conditions Tested section of the report. The region of interest (ROI) of the test comprises these regions, in addition to the 20 intronic bases flanking each exon. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to high coverage and the sequences are compared to standards and references of normal variation. More than 99% of all bases in the ROI are sequenced at greater than the minimum read depth. Mutations may not be detected in areas of lower sequence coverage. Small insertions and deletions may not be as accurately determined as single nucleotide variants. Genes that have closely related pseudogenes may be addressed by a different method. *CFTR* and *DMD* testing includes analysis for both large (exon-level) deletions and duplications with an average sensitivity of 99%, while other genes are only analyzed for large deletions with a sensitivity of >75%. However, the sensitivity may be higher for selected founder deletions. If *GJB2* is tested, two large upstream deletions which overlap *GJB6* and affect the expression of *GJB2*, del(*GJB6*-D13S1830) and del(*GJB6*-D13S1854), are also analyzed. Mosaicism or somatic variants present at low levels may not be detected. If detected, these may not be reported.

Detection rates are determined by using literature to estimate the fraction of disease alleles, weighted by frequency, that the methodology is unable to detect. Detection rates only account for analytical sensitivity and certain variants that have been previously described in the literature may not be reported if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease-specific rates of de novo mutations.

All variants that are a recognized cause of the disease will be reported. In addition, variants that have not previously been established as a recognized cause of disease may be identified. In these cases, only variants classified as "likely" pathogenic are reported. Likely pathogenic variants are described elsewhere in the report as "likely to have a negative impact on gene function". Likely pathogenic variants are evaluated and classified by assessing the nature of the variant and reviewing reports of allele frequencies in cases and controls, functional studies, variant annotation and effect prediction, and segregation studies. Exon level duplications are assumed to be in tandem and are classified according to their predicted effect on the reading frame. Benign variants, variants of uncertain significance, and variants not directly associated with the intended disease phenotype are not reported. Curation summaries of reported variants are available upon request.

Spinal muscular atrophy

Targeted copy number analysis is used to determine the copy number of exon 7 of the *SMN1* gene relative to other genes. Other mutations may interfere with this analysis. Some individuals with two copies of *SMN1* are carriers with two *SMN1* genes on one chromosome and a *SMN1* deletion on the other chromosome. This is more likely in individuals who have 2 copies of the *SMN1* gene and are positive for the g.27134T>G SNP, which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have 2 copies of *SMN1*.

Analysis of homologous regions

A combination of high-throughput sequencing, read depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss of function mutations in certain genes that have homology to other regions. The precise breakpoints of large deletions in these genes cannot be determined, but are estimated from copy number analysis. High numbers of pseudogene copies may interfere with this analysis.

If *CYP21A2* is tested, patients who have one or more additional copies of the *CYP21A2* gene and a loss of function mutation may not actually be a carrier of 21-hydroxylase-deficient congenital adrenal hyperplasia (CAH). Because the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are only based on published incidences for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate of overall chances for 21-hydroxylase-deficient CAH, especially in the aforementioned populations, as they do not account for non-classic CAH. If *HBA1*/*HBA2* are tested, some individuals with four alpha globin genes may be carriers, with three genes on one chromosome and a deletion on the other chromosome. This and similar, but rare, carrier states, where complementary changes exist in both the gene and a pseudogene, may not be detected by the assay.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Limitations

In an unknown number of cases, nearby genetic variants may interfere with mutation detection. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions and technical errors. This test is designed to detect and report germline alterations. While somatic variants present at low levels may be detected, these may not be reported. If more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes. The test does not fully address all inherited forms of intellectual disability, birth defects and genetic disease. A family history of any of these conditions may warrant additional evaluation. Furthermore, not all mutations will be identified in the genes analyzed and additional testing may be beneficial for some patients. For example, individuals of African, Southeast Asian, and Mediterranean ancestry are at increased risk for being carriers for hemoglobinopathies, which can be identified by CBC and hemoglobin electrophoresis or HPLC (*ACOG Practice Bulletin No. 78. Obstet.Gynecol. 2007;109:229-37*).

This test was developed and its performance characteristics determined by Myriad Women's Health, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: **#05D1102604**.

LABORATORY DIRECTOR

Hyunseok Kang

H. Peter Kang, MD, MS, FCAP

Report content approved by Saurav Guha, PhD, FACMG on Sep 10, 2018

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Conditions Tested

11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia - **Gene:** CYP11B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000497:1-9. **Detection Rate:** Southeast Asian 94%.
21-hydroxylase-deficient Congenital Adrenal Hyperplasia - **Gene:** CYP21A2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs*21, I173N, L308Ffs*6, P31L, Q319*, Q319*+CYP21A2dup, R357W, V281L, [I237N;V238E;M240K], c.293-13C>G. **Detection Rate:** Southeast Asian 88%.
6-pyruvoyl-tetrahydropterin Synthase Deficiency - **Gene:** PTS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000317:1-6. **Detection Rate:** Southeast Asian >99%.
ABCC8-related Hyperinsulinism - **Gene:** ABCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000352:1-39. **Detection Rate:** Southeast Asian >99%.
Adenosine Deaminase Deficiency - **Gene:** ADA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000022:1-12. **Detection Rate:** Southeast Asian >99%.
Alpha Thalassemia - **Genes:** HBA1, HBA2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** -(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, anti3.7, anti4.2, del HS-40. **Detection Rate:** Southeast Asian 90%.
Alpha-mannosidosis - **Gene:** MAN2B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000528:1-23. **Detection Rate:** Southeast Asian >99%.
Alpha-sarcoglycanopathy - **Gene:** SGCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000023:1-9. **Detection Rate:** Southeast Asian >99%.
Alstrom Syndrome - **Gene:** ALMS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015120:1-23. **Detection Rate:** Southeast Asian >99%.
AMT-related Glycine Encephalopathy - **Gene:** AMT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000481:1-9. **Detection Rate:** Southeast Asian >99%.
Andermann Syndrome - **Gene:** SLC12A6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133647:1-25. **Detection Rate:** Southeast Asian >99%.
Argininemia - **Gene:** ARG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001244438:1-8. **Detection Rate:** Southeast Asian 97%.
Argininosuccinic Aciduria - **Gene:** ASL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001024943:1-16. **Detection Rate:** Southeast Asian >99%.
ARSACS - **Gene:** SACS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014363:2-10. **Detection Rate:** Southeast Asian 99%.
Aspartylglycosaminuria - **Gene:** AGA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000027:1-9. **Detection Rate:** Southeast Asian >99%.
Ataxia with Vitamin E Deficiency - **Gene:** TTPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000370:1-5. **Detection Rate:** Southeast Asian >99%.
Ataxia-telangiectasia - **Gene:** ATM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000051:2-63. **Detection Rate:** Southeast Asian >99%.
ATP7A-related Disorders - **Gene:** ATP7A. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000052:2-23. **Detection Rate:** Southeast Asian 92%.
Autosomal Recessive Osteopetrosis Type 1 - **Gene:** TCIRG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006019:2-20. **Detection Rate:** Southeast Asian >99%.
Bardet-Biedl Syndrome, BBS1-related - **Gene:** BBS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024649:1-17. **Detection Rate:** Southeast Asian >99%.
Bardet-Biedl Syndrome, BBS10-related - **Gene:** BBS10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024685:1-2. **Detection Rate:** Southeast Asian >99%.
Bardet-Biedl Syndrome, BBS12-related - **Gene:** BBS12. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_152618:2. **Detection Rate:** Southeast Asian >99%.
Bardet-Biedl Syndrome, BBS2-related - **Gene:** BBS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_031885:1-17. **Detection Rate:** Southeast Asian >99%.
Beta-sarcoglycanopathy - **Gene:** SGCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000232:1-6. **Detection Rate:** Southeast Asian >99%.
Biotinidase Deficiency - **Gene:** BTD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000060:1-4. **Detection Rate:** Southeast Asian >99%.
Bloom Syndrome - **Gene:** BLM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000057:2-22. **Detection Rate:** Southeast Asian >99%.
Calpainopathy - **Gene:** CAPN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000070:1-24. **Detection Rate:** Southeast Asian >99%.
Canavan Disease - **Gene:** ASPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000049:1-6. **Detection Rate:** Southeast Asian 98%.
Carbamoylphosphate Synthetase I Deficiency - **Gene:** CPS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001875:1-38. **Detection Rate:** Southeast Asian >99%.
Carnitine Palmitoyltransferase IA Deficiency - **Gene:** CPT1A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001876:2-19. **Detection Rate:** Southeast Asian >99%.
Carnitine Palmitoyltransferase II Deficiency - **Gene:** CPT2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000098:1-5. **Detection Rate:** Southeast Asian >99%.
Cartilage-hair Hypoplasia - **Gene:** RMRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NR_003051:1. **Detection Rate:** Southeast Asian >99%.
Cerebrotendinous Xanthomatosis - **Gene:** CYP27A1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000784:1-9. **Detection Rate:** Southeast Asian >99%.
Citrullinemia Type 1 - **Gene:** ASS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000050:3-16. **Detection Rate:** Southeast Asian >99%.
CLN3-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001042432:2-16. **Detection Rate:** Southeast Asian >99%.
CLN5-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006493:1-4. **Detection Rate:** Southeast Asian >99%.
CLN6-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017882:1-7. **Detection Rate:** Southeast Asian >99%.
Cohen Syndrome - **Gene:** VPS13B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017890:2-62. **Detection Rate:** Southeast Asian 97%.
COL4A3-related Alport Syndrome - **Gene:** COL4A3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000091:1-52. **Detection Rate:** Southeast Asian 97%.
COL4A4-related Alport Syndrome - **Gene:** COL4A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000092:2-48. **Detection Rate:** Southeast Asian 98%.
Congenital Disorder of Glycosylation Type Ia - **Gene:** PMM2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000303:1-8. **Detection Rate:** Southeast Asian >99%.
Congenital Disorder of Glycosylation Type Ib - **Gene:** MPI. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002435:1-8. **Detection Rate:** Southeast Asian >99%.
Congenital Disorder of Glycosylation Type Ic - **Gene:** ALG6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_013339:2-15. **Detection Rate:** Southeast Asian >99%.
Congenital Finnish Nephrosis - **Gene:** NPHS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004646:1-29. **Detection Rate:** Southeast Asian >99%.
Costeff Optic Atrophy Syndrome - **Gene:** OPA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_025136:1-2. **Detection Rate:** Southeast Asian >99%.
Cystic Fibrosis - **Gene:** CFTR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. **Detection Rate:** Southeast Asian >99%.
Cystinosis - **Gene:** CTNS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004937:3-12. **Detection Rate:** Southeast Asian >99%.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

D-bifunctional Protein Deficiency - **Gene:** HSD17B4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000414:1-24. **Detection Rate:** Southeast Asian 98%.
Delta-sarcoglycanopathy - **Gene:** SGCD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000337:2-9. **Detection Rate:** Southeast Asian 99%.
Dysferlinopathy - **Gene:** DYSF. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001130987:1-56. **Detection Rate:** Southeast Asian 98%.
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy) - **Gene:** DMD. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_004006:1-79. **Detection Rate:** Southeast Asian >99%.
ERCC6-related Disorders - **Gene:** ERCC6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000124:2-21. **Detection Rate:** Southeast Asian 99%.
ERCC8-related Disorders - **Gene:** ERCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000082:1-12. **Detection Rate:** Southeast Asian 95%.
EVC-related Ellis-van Creveld Syndrome - **Gene:** EVC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153717:1-21. **Detection Rate:** Southeast Asian 96%.
EVC2-related Ellis-van Creveld Syndrome - **Gene:** EVC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_147127:1-22. **Detection Rate:** Southeast Asian >99%.
Fabry Disease - **Gene:** GLA. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000169:1-7. **Detection Rate:** Southeast Asian 98%.
Familial Dysautonomia - **Gene:** IKBKAP. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003640:2-37. **Detection Rate:** Southeast Asian >99%.
Familial Mediterranean Fever - **Gene:** MEFV. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000243:1-10. **Detection Rate:** Southeast Asian >99%.
Fanconi Anemia Complementation Group A - **Gene:** FANCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000135:1-43. **Detection Rate:** Southeast Asian 92%.
Fanconi Anemia Type C - **Gene:** FANCC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000136:2-15. **Detection Rate:** Southeast Asian >99%.
FKRP-related Disorders - **Gene:** FKRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_024301:4. **Detection Rate:** Southeast Asian >99%.
FKTN-related Disorders - **Gene:** FKTN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001079802:3-11. **Detection Rate:** Southeast Asian >99%.
Galactokinase Deficiency - **Gene:** GALK1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000154:1-8. **Detection Rate:** Southeast Asian >99%.
Galactosemia - **Gene:** GALT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000155:1-11. **Detection Rate:** Southeast Asian >99%.
Gamma-sarcoglycanopathy - **Gene:** SGCG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000231:2-8. **Detection Rate:** Southeast Asian 88%.
Gaucher Disease - **Gene:** GBA. Autosomal Recessive. Analysis of homologous regions. **Variants (10):** D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs*18. **Detection Rate:** Southeast Asian 60%.
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness - **Gene:** GJB2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004004:1-2. **Detection Rate:** Southeast Asian >99%.
GLB1-related Disorders - **Gene:** GLB1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000404:1-16. **Detection Rate:** Southeast Asian >99%.
GLDC-related Glycine Encephalopathy - **Gene:** GLDC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000170:1-25. **Detection Rate:** Southeast Asian 94%.
Glutaric Acidemia Type 1 - **Gene:** GCDH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000159:2-12. **Detection Rate:** Southeast Asian >99%.
Glycogen Storage Disease Type Ia - **Gene:** G6PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000151:1-5. **Detection Rate:** Southeast Asian >99%.
Glycogen Storage Disease Type Ib - **Gene:** SLC37A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001164277:3-11. **Detection Rate:** Southeast Asian >99%.
Glycogen Storage Disease Type III - **Gene:** AGL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000642:2-34. **Detection Rate:** Southeast Asian >99%.
GNPTAB-related Disorders - **Gene:** GNPTAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024312:1-21. **Detection Rate:** Southeast Asian >99%.
GRACILE Syndrome - **Gene:** BCS1L. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004328:3-9. **Detection Rate:** Southeast Asian >99%.
HADHA-related Disorders - **Gene:** HADHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000182:1-20. **Detection Rate:** Southeast Asian >99%.
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease) - **Gene:** HBB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000518:1-3. **Detection Rate:** Southeast Asian >99%.
Hereditary Fructose Intolerance - **Gene:** ALDOB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000035:2-9. **Detection Rate:** Southeast Asian >99%.
Herlitz Junctional Epidermolysis Bullosa, LAMA3-related - **Gene:** LAMA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000227:1-38. **Detection Rate:** Southeast Asian >99%.
Herlitz Junctional Epidermolysis Bullosa, LAMB3-related - **Gene:** LAMB3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000228:2-23. **Detection Rate:** Southeast Asian >99%.
Herlitz Junctional Epidermolysis Bullosa, LAMC2-related - **Gene:** LAMC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005562:1-23. **Detection Rate:** Southeast Asian >99%.
Hexosaminidase A Deficiency (Including Tay-Sachs Disease) - **Gene:** HEXA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000520:1-14. **Detection Rate:** Southeast Asian >99%.
HMG-CoA Lyase Deficiency - **Gene:** HMGCL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000191:1-9. **Detection Rate:** Southeast Asian 98%.
Holocarboxylase Synthetase Deficiency - **Gene:** HLCS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000411:4-12. **Detection Rate:** Southeast Asian >99%.
Homocystinuria Caused by Cystathionine Beta-synthase Deficiency - **Gene:** CBS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000071:3-17. **Detection Rate:** Southeast Asian >99%.
Hydrolethalus Syndrome - **Gene:** HYLS1. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_001134793:3. **Detection Rate:** Southeast Asian >99%.
Hypophosphatasia, Autosomal Recessive - **Gene:** ALPL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000478:2-12. **Detection Rate:** Southeast Asian >99%.
Inclusion Body Myopathy 2 - **Gene:** GNE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001128227:1-12. **Detection Rate:** Southeast Asian >99%.
Isovaleric Acidemia - **Gene:** IVD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002225:1-12. **Detection Rate:** Southeast Asian >99%.
Joubert Syndrome 2 - **Gene:** TMEM216. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001173990:1-5. **Detection Rate:** Southeast Asian >99%.
KCNJ11-related Familial Hyperinsulinism - **Gene:** KCNJ11. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000525:1. **Detection Rate:** Southeast Asian >99%.
Krabbe Disease - **Gene:** GALC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000153:1-17. **Detection Rate:** Southeast Asian >99%.
LAMA2-related Muscular Dystrophy - **Gene:** LAMA2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000426:1-65. **Detection Rate:** Southeast Asian >99%.
Leigh Syndrome, French-Canadian Type - **Gene:** LRPPRC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133259:1-38. **Detection Rate:** Southeast Asian >99%.
Lipoamide Dehydrogenase Deficiency - **Gene:** DLD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000108:1-14. **Detection Rate:** Southeast Asian >99%.
Lipoid Congenital Adrenal Hyperplasia - **Gene:** STAR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000349:1-7. **Detection Rate:** Southeast Asian >99%.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Lysosomal Acid Lipase Deficiency - **Gene:** LIPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000235:2-10. **Detection Rate:** Southeast Asian >99%.
Maple Syrup Urine Disease Type 1B - **Gene:** BCKDHB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_183050:1-10. **Detection Rate:** Southeast Asian >99%.
Maple Syrup Urine Disease Type Ia - **Gene:** BCKDHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000709:1-9. **Detection Rate:** Southeast Asian >99%.
Maple Syrup Urine Disease Type II - **Gene:** DBT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001918:1-11. **Detection Rate:** Southeast Asian 96%.
Medium Chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000016:1-12. **Detection Rate:** Southeast Asian >99%.
Megalencephalic Leukoencephalopathy with Subcortical Cysts - **Gene:** MLC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015166:2-12. **Detection Rate:** Southeast Asian >99%.
Metachromatic Leukodystrophy - **Gene:** ARSA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000487:1-8. **Detection Rate:** Southeast Asian >99%.
Methylmalonic Acidemia, cblA Type - **Gene:** MMAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_172250:2-7. **Detection Rate:** Southeast Asian >99%.
Methylmalonic Acidemia, cblB Type - **Gene:** MMAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_052845:1-9. **Detection Rate:** Southeast Asian >99%.
Methylmalonic Aciduria and Homocystinuria, cblC Type - **Gene:** MMACHC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015506:1-4. **Detection Rate:** Southeast Asian >99%.
MKS1-related Disorders - **Gene:** MKS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017777:1-18. **Detection Rate:** Southeast Asian >99%.
Mucolipidosis III Gamma - **Gene:** GNPTG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032520:1-11. **Detection Rate:** Southeast Asian >99%.
Mucolipidosis IV - **Gene:** MCOLN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_020533:1-14. **Detection Rate:** Southeast Asian >99%.
Mucopolysaccharidosis Type I - **Gene:** IDUA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000203:1-14. **Detection Rate:** Southeast Asian >99%.
Mucopolysaccharidosis Type II - **Gene:** IDS. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000202:1-9. **Detection Rate:** Southeast Asian 88%.
Mucopolysaccharidosis Type IIIA - **Gene:** SGSH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000199:1-8. **Detection Rate:** Southeast Asian >99%.
Mucopolysaccharidosis Type IIIB - **Gene:** NAGLU. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000263:1-6. **Detection Rate:** Southeast Asian >99%.
Mucopolysaccharidosis Type IIIC - **Gene:** HGSNAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_152419:1-18. **Detection Rate:** Southeast Asian >99%.
Muscle-eye-brain Disease - **Gene:** POMGNT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017739:2-22. **Detection Rate:** Southeast Asian 96%.
MUT-related Methylmalonic Acidemia - **Gene:** MUT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000255:2-13. **Detection Rate:** Southeast Asian >99%.
MYO7A-related Disorders - **Gene:** MYO7A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000260:2-49. **Detection Rate:** Southeast Asian >99%.
NEB-related Nemaline Myopathy - **Gene:** NEB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001271208:3-80,117-183. **Detection Rate:** Southeast Asian 92%.
Nephrotic Syndrome, NPHS2-related - **Gene:** NPHS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014625:1-8. **Detection Rate:** Southeast Asian >99%.
Niemann-Pick Disease Type C - **Gene:** NPC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000271:1-25. **Detection Rate:** Southeast Asian >99%.
Niemann-Pick Disease Type C2 - **Gene:** NPC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006432:1-5. **Detection Rate:** Southeast Asian >99%.
Niemann-Pick Disease, SMPD1-associated - **Gene:** SMPD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000543:1-6. **Detection Rate:** Southeast Asian >99%.
Nijmegen Breakage Syndrome - **Gene:** NBN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002485:1-16. **Detection Rate:** Southeast Asian >99%.
Northern Epilepsy - **Gene:** CLN8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_018941:2-3. **Detection Rate:** Southeast Asian >99%.
Ornithine Transcarbamylase Deficiency - **Gene:** OTC. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000531:1-10. **Detection Rate:** Southeast Asian 97%.
PCCA-related Propionic Acidemia - **Gene:** PCCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000282:1-24. **Detection Rate:** Southeast Asian 95%.
PCCB-related Propionic Acidemia - **Gene:** PCCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001178014:1-16. **Detection Rate:** Southeast Asian >99%.
PCDH15-related Disorders - **Gene:** PCDH15. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_033056:2-33. **Detection Rate:** Southeast Asian 93%.
Pendred Syndrome - **Gene:** SLC26A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000441:2-21. **Detection Rate:** Southeast Asian >99%.
Peroxisome Biogenesis Disorder Type 3 - **Gene:** PEX12. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000286:1-3. **Detection Rate:** Southeast Asian >99%.
Peroxisome Biogenesis Disorder Type 4 - **Gene:** PEX6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000287:1-17. **Detection Rate:** Southeast Asian 97%.
Peroxisome Biogenesis Disorder Type 5 - **Gene:** PEX2. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000318:4. **Detection Rate:** Southeast Asian >99%.
Peroxisome Biogenesis Disorder Type 6 - **Gene:** PEX10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153818:1-6. **Detection Rate:** Southeast Asian >99%.
PEX1-related Zellweger Syndrome Spectrum - **Gene:** PEX1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000466:1-24. **Detection Rate:** Southeast Asian >99%.
Phenylalanine Hydroxylase Deficiency - **Gene:** PAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000277:1-13. **Detection Rate:** Southeast Asian >99%.
PKHD1-related Autosomal Recessive Polycystic Kidney Disease - **Gene:** PKHD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138694:2-67. **Detection Rate:** Southeast Asian >99%.
Polyglandular Autoimmune Syndrome Type 1 - **Gene:** AIRE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000383:1-14. **Detection Rate:** Southeast Asian >99%.
Pompe Disease - **Gene:** GAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000152:2-20. **Detection Rate:** Southeast Asian >99%.
PPT1-related Neuronal Ceroid Lipofuscinosis - **Gene:** PPT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000310:1-9. **Detection Rate:** Southeast Asian >99%.
Primary Carnitine Deficiency - **Gene:** SLC22A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003060:1-10. **Detection Rate:** Southeast Asian >99%.
Primary Hyperoxaluria Type 1 - **Gene:** AGXT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000030:1-11. **Detection Rate:** Southeast Asian >99%.
Primary Hyperoxaluria Type 2 - **Gene:** GRHPR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012203:1-9. **Detection Rate:** Southeast Asian >99%.
Primary Hyperoxaluria Type 3 - **Gene:** HOGA1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138413:1-7. **Detection Rate:** Southeast Asian >99%.
PROP1-related Combined Pituitary Hormone Deficiency - **Gene:** PROP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006261:1-3. **Detection Rate:** Southeast Asian >99%.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Pycnodysostosis - **Gene:** CTSK. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000396:2-8. **Detection Rate:** Southeast Asian >99%.
Pyruvate Carboxylase Deficiency - **Gene:** PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_022172:2-21. **Detection Rate:** Southeast Asian >99%.
Rhizomelic Chondrodysplasia Punctata Type 1 - **Gene:** PEX7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000288:1-10. **Detection Rate:** Southeast Asian >99%.
RTEL1-related Disorders - **Gene:** RTEL1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032957:2-35. **Detection Rate:** Southeast Asian >99%.
Salla Disease - **Gene:** SLC17A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012434:1-11. **Detection Rate:** Southeast Asian 98%.
Sandhoff Disease - **Gene:** HEXB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000521:1-14. **Detection Rate:** Southeast Asian 99%.
Segawa Syndrome - **Gene:** TH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000360:1-13. **Detection Rate:** Southeast Asian >99%.
Short Chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000017:1-10. **Detection Rate:** Southeast Asian >99%.
Sjogren-Larsson Syndrome - **Gene:** ALDH3A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000382:1-10. **Detection Rate:** Southeast Asian 97%.
Smith-Lemli-Opitz Syndrome - **Gene:** DHCR7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001360:3-9. **Detection Rate:** Southeast Asian >99%.
Spastic Paraplegia Type 15 - **Gene:** ZFYVE26. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015346:2-42. **Detection Rate:** Southeast Asian >99%.
Spinal Muscular Atrophy - **Gene:** SMN1. Autosomal Recessive. Spinal muscular atrophy. **Variant (1):** SMN1 copy number. **Detection Rate:** Southeast Asian 93%.
Spondylothoracic Dysostosis - **Gene:** MESP2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001039958:1-2. **Detection Rate:** Southeast Asian >99%.
Sulfate Transporter-related Osteochondrodysplasia - **Gene:** SLC26A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000112:2-3. **Detection Rate:** Southeast Asian >99%.
TGM1-related Autosomal Recessive Congenital Ichthyosis - **Gene:** TGM1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000359:2-15. **Detection Rate:** Southeast Asian >99%.
TPP1-related Neuronal Ceroid Lipofuscinosis - **Gene:** TPP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000391:1-13. **Detection Rate:** Southeast Asian >99%.
Tyrosinemia Type I - **Gene:** FAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000137:1-14. **Detection Rate:** Southeast Asian >99%.
Tyrosinemia Type II - **Gene:** TAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000353:2-12. **Detection Rate:** Southeast Asian >99%.
USH1C-related Disorders - **Gene:** USH1C. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153676:1-27. **Detection Rate:** Southeast Asian >99%.
USH2A-related Disorders - **Gene:** USH2A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_206933:2-72. **Detection Rate:** Southeast Asian 94%.
Usher Syndrome Type 3 - **Gene:** CLRN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_174878:1-3. **Detection Rate:** Southeast Asian >99%.
Very Long Chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADVL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000018:1-20. **Detection Rate:** Southeast Asian >99%.
Wilson Disease - **Gene:** ATP7B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000053:1-21. **Detection Rate:** Southeast Asian >99%.
X-linked Adrenoleukodystrophy - **Gene:** ABCD1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000033:1-6. **Detection Rate:** Southeast Asian 77%.
X-linked Alport Syndrome - **Gene:** COL4A5. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000495:1-51. **Detection Rate:** Southeast Asian 95%.
X-linked Congenital Adrenal Hypoplasia - **Gene:** NR0B1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000475:1-2. **Detection Rate:** Southeast Asian 99%.
X-linked Juvenile Retinoschisis - **Gene:** RS1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000330:1-6. **Detection Rate:** Southeast Asian 98%.
X-linked Myotubular Myopathy - **Gene:** MTM1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000252:2-15. **Detection Rate:** Southeast Asian 98%.
X-linked Severe Combined Immunodeficiency - **Gene:** IL2RG. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000206:1-8. **Detection Rate:** Southeast Asian >99%.
Xeroderma Pigmentosum Group A - **Gene:** XPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000380:1-6. **Detection Rate:** Southeast Asian >99%.
Xeroderma Pigmentosum Group C - **Gene:** XPC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004628:1-16. **Detection Rate:** Southeast Asian 97%.

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Risk Calculations

Below are the risk calculations for all conditions tested. Since negative results do not completely rule out the possibility of being a carrier, the **residual risk** represents the patient's post-test likelihood of being a carrier and the **reproductive risk** represents the likelihood the patient's future children could inherit each disease. These risks are inherent to all carrier screening tests, may vary by ethnicity, are predicated on a negative family history and are present even after a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

†Indicates a positive result. See the full clinical report for interpretation and details.

Disease	DONOR 10289 Residual Risk	Reproductive Risk
11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia	1 in 3,300	< 1 in 1,000,000
21-hydroxylase-deficient Congenital Adrenal Hyperplasia	1 in 480	1 in 110,000
6-pyruvoyl-tetrahydropterin Synthase Deficiency	< 1 in 50,000	< 1 in 1,000,000
ABCC8-related Hyperinsulinism	1 in 11,000	< 1 in 1,000,000
Adenosine Deaminase Deficiency	1 in 39,000	< 1 in 1,000,000
Alpha Thalassemia	Alpha globin status: aa/aa.	Not calculated
Alpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Alpha-sarcoglycanopathy	1 in 45,000	< 1 in 1,000,000
Alstrom Syndrome	< 1 in 50,000	< 1 in 1,000,000
AMT-related Glycine Encephalopathy	1 in 22,000	< 1 in 1,000,000
Andermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
Argininemia	< 1 in 17,000	< 1 in 1,000,000
Argininosuccinic Aciduria	1 in 13,000	< 1 in 1,000,000
ARSACS	< 1 in 44,000	< 1 in 1,000,000
Aspartylglycosaminuria	< 1 in 50,000	< 1 in 1,000,000
Ataxia with Vitamin E Deficiency	< 1 in 50,000	< 1 in 1,000,000
Ataxia-telangiectasia	1 in 16,000	< 1 in 1,000,000
ATP7A-related Disorders	< 1 in 1,000,000	1 in 600,000
Autosomal Recessive Osteopetrosis Type 1	1 in 35,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS1-related	1 in 16,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS10-related	1 in 16,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS12-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS2-related	< 1 in 50,000	< 1 in 1,000,000
Beta-sarcoglycanopathy	NM_000232.4(SGCB):c.1A>G(M1?) heterozygote †	1 in 2,000
Biotinidase Deficiency	1 in 18,000	< 1 in 1,000,000
Bloom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Calpainopathy	1 in 13,000	< 1 in 1,000,000
Canavan Disease	< 1 in 31,000	< 1 in 1,000,000
Carbamoylphosphate Synthetase I Deficiency	< 1 in 57,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase II Deficiency	< 1 in 50,000	< 1 in 1,000,000
Cartilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
Cerebrotendinous Xanthomatosis	1 in 11,000	< 1 in 1,000,000
Citrullinemia Type 1	1 in 12,000	< 1 in 1,000,000
CLN3-related Neuronal Ceroid Lipofuscinosis	1 in 22,000	< 1 in 1,000,000
CLN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN6-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Cohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
COL4A3-related Alport Syndrome	1 in 11,000	< 1 in 1,000,000
COL4A4-related Alport Syndrome	1 in 21,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ia	1 in 16,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ib	< 1 in 50,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ic	< 1 in 50,000	< 1 in 1,000,000
Congenital Finnish Nephrosis	< 1 in 50,000	< 1 in 1,000,000
Costeff Optic Atrophy Syndrome	< 1 in 50,000	< 1 in 1,000,000
Cystic Fibrosis	1 in 8,600	< 1 in 1,000,000
Cystinosis	1 in 22,000	< 1 in 1,000,000
D-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 10289 Residual Risk	Reproductive Risk
Delta-sarcoglycanopathy	< 1 in 40,000	< 1 in 1,000,000
Dysferlinopathy	1 in 11,000	< 1 in 1,000,000
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)	Not calculated	Not calculated
ERCC6-related Disorders	1 in 19,000	< 1 in 1,000,000
ERCC8-related Disorders	1 in 7,300	< 1 in 1,000,000
EVC-related Ellis-van Creveld Syndrome	1 in 7,500	< 1 in 1,000,000
EVC2-related Ellis-van Creveld Syndrome	< 1 in 50,000	< 1 in 1,000,000
Fabry Disease	< 1 in 1,000,000	1 in 80,000
Familial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
Familial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
Fanconi Anemia Complementation Group A	1 in 3,100	< 1 in 1,000,000
Fanconi Anemia Type C	1 in 16,000	< 1 in 1,000,000
FKRP-related Disorders	< 1 in 50,000	< 1 in 1,000,000
FKTN-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Galactokinase Deficiency	1 in 35,000	< 1 in 1,000,000
Galactosemia	< 1 in 50,000	< 1 in 1,000,000
Gamma-sarcoglycanopathy	1 in 3,000	< 1 in 1,000,000
Gaucher Disease	1 in 310	1 in 150,000
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 10,000	< 1 in 1,000,000
GLB1-related Disorders	1 in 19,000	< 1 in 1,000,000
GLDC-related Glycine Encephalopathy	1 in 2,800	< 1 in 1,000,000
Glutaric Acidemia Type 1	1 in 10,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ia	1 in 18,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ib	1 in 35,000	< 1 in 1,000,000
Glycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
GNPTAB-related Disorders	1 in 32,000	< 1 in 1,000,000
GRACILE Syndrome	< 1 in 50,000	< 1 in 1,000,000
HADHA-related Disorders	1 in 15,000	< 1 in 1,000,000
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease)	1 in 2,400	1 in 240,000
Hereditary Fructose Intolerance	< 1 in 50,000	< 1 in 1,000,000
Herlitz Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
Herlitz Junctional Epidermolysis Bullosa, LAMB3-related	< 1 in 50,000	< 1 in 1,000,000
Herlitz Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
Hexosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
HMG-CoA Lyase Deficiency	< 1 in 33,000	< 1 in 1,000,000
Holocarboxylase Synthetase Deficiency	1 in 15,000	< 1 in 1,000,000
Homocystinuria Caused by Cystathionine Beta-synthase Deficiency	1 in 25,000	< 1 in 1,000,000
Hydrolethalus Syndrome	< 1 in 50,000	< 1 in 1,000,000
Hypophosphatasia, Autosomal Recessive	1 in 16,000	< 1 in 1,000,000
Inclusion Body Myopathy 2	< 1 in 50,000	< 1 in 1,000,000
Isovaleric Acidemia	1 in 25,000	< 1 in 1,000,000
Joubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
KCNJ11-related Familial Hyperinsulinism	< 1 in 50,000	< 1 in 1,000,000
Krabbe Disease	1 in 15,000	< 1 in 1,000,000
LAMA2-related Muscular Dystrophy	1 in 17,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type	< 1 in 50,000	< 1 in 1,000,000
Lipoamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Lipoid Congenital Adrenal Hyperplasia	< 1 in 50,000	< 1 in 1,000,000
Lysosomal Acid Lipase Deficiency	1 in 30,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type 1B	1 in 25,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ia	1 in 19,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type II	1 in 7,600	< 1 in 1,000,000
Medium Chain Acyl-CoA Dehydrogenase Deficiency	1 in 11,000	< 1 in 1,000,000
Megalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000	< 1 in 1,000,000
Metachromatic Leukodystrophy	1 in 20,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblA Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblB Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Aciduria and Homocystinuria, cblC Type	1 in 16,000	< 1 in 1,000,000
MKS1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis III Gamma	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis IV	< 1 in 50,000	< 1 in 1,000,000

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 10289 Residual Risk	Reproductive Risk
Mucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type II	< 1 in 1,000,000	1 in 300,000
Mucopolysaccharidosis Type IIIA	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIB	1 in 31,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIC	1 in 43,000	< 1 in 1,000,000
Muscle-eye-brain Disease	< 1 in 12,000	< 1 in 1,000,000
MUT-related Methylmalonic Acidemia	1 in 5,300	< 1 in 1,000,000
MYO7A-related Disorders	1 in 15,000	< 1 in 1,000,000
NEB-related Nemaline Myopathy	< 1 in 6,700	< 1 in 1,000,000
Nephrotic Syndrome, NPHS2-related	1 in 35,000	< 1 in 1,000,000
Niemann-Pick Disease Type C	1 in 19,000	< 1 in 1,000,000
Niemann-Pick Disease Type C2	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease, SMPD1-associated	1 in 25,000	< 1 in 1,000,000
Nijmegen Breakage Syndrome	1 in 16,000	< 1 in 1,000,000
Northern Epilepsy	< 1 in 50,000	< 1 in 1,000,000
Ornithine Transcarbamylase Deficiency	< 1 in 1,000,000	1 in 140,000
PCCA-related Propionic Acidemia	1 in 4,200	< 1 in 1,000,000
PCCB-related Propionic Acidemia	1 in 22,000	< 1 in 1,000,000
PCDH15-related Disorders	1 in 5,300	< 1 in 1,000,000
Pendred Syndrome	1 in 7,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 3	1 in 44,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 4	1 in 9,300	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 5	< 1 in 71,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 6	< 1 in 50,000	< 1 in 1,000,000
PEX1-related Zellweger Syndrome Spectrum	1 in 35,000	< 1 in 1,000,000
Phenylalanine Hydroxylase Deficiency	1 in 5,000	1 in 990,000
PKHD1-related Autosomal Recessive Polycystic Kidney Disease	< 1 in 50,000	< 1 in 1,000,000
Polyglandular Autoimmune Syndrome Type 1	< 1 in 50,000	< 1 in 1,000,000
Pompe Disease	1 in 11,000	< 1 in 1,000,000
PPT1-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Primary Carnitine Deficiency	1 in 16,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 1	1 in 35,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 3	1 in 20,000	< 1 in 1,000,000
PROP1-related Combined Pituitary Hormone Deficiency	1 in 11,000	< 1 in 1,000,000
Pycnodysostosis	< 1 in 50,000	< 1 in 1,000,000
Pyruvate Carboxylase Deficiency	1 in 25,000	< 1 in 1,000,000
Rhizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
RTEL1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Salla Disease	< 1 in 30,000	< 1 in 1,000,000
Sandhoff Disease	1 in 30,000	< 1 in 1,000,000
Segawa Syndrome	< 1 in 50,000	< 1 in 1,000,000
Short Chain Acyl-CoA Dehydrogenase Deficiency	1 in 16,000	< 1 in 1,000,000
Sjogren-Larsson Syndrome	1 in 9,100	< 1 in 1,000,000
Smith-Lemli-Opitz Syndrome	< 1 in 50,000	< 1 in 1,000,000
Spastic Paraplegia Type 15	< 1 in 50,000	< 1 in 1,000,000
Spinal Muscular Atrophy	Negative for g.27134T>G SNP SMN1: 2 copies 1 in 700	1 in 150,000
Spondylothoracic Dysostosis	< 1 in 50,000	< 1 in 1,000,000
Sulfate Transporter-related Osteochondrodysplasia	1 in 11,000	< 1 in 1,000,000
TGM1-related Autosomal Recessive Congenital Ichthyosis	1 in 22,000	< 1 in 1,000,000
TPP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
Tyrosinemia Type I	1 in 17,000	< 1 in 1,000,000
Tyrosinemia Type II	1 in 25,000	< 1 in 1,000,000
USH1C-related Disorders	1 in 35,000	< 1 in 1,000,000
USH2A-related Disorders	1 in 2,200	< 1 in 1,000,000
Usher Syndrome Type 3	< 1 in 50,000	< 1 in 1,000,000
Very Long Chain Acyl-CoA Dehydrogenase Deficiency	1 in 8,800	< 1 in 1,000,000
Wilson Disease	1 in 5,000	1 in 990,000
X-linked Adrenoleukodystrophy	1 in 90,000	1 in 42,000
X-linked Alport Syndrome	Not calculated	Not calculated

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 10289 Residual Risk	Reproductive Risk
X-linked Congenital Adrenal Hypoplasia	< 1 in 1,000,000	< 1 in 1,000,000
X-linked Juvenile Retinoschisis	< 1 in 1,000,000	1 in 50,000
X-linked Myotubular Myopathy	Not calculated	Not calculated
X-linked Severe Combined Immunodeficiency	< 1 in 1,000,000	1 in 200,000
Xeroderma Pigmentosum Group A	< 1 in 50,000	< 1 in 1,000,000
Xeroderma Pigmentosum Group C	1 in 7,300	< 1 in 1,000,000

Counsyl is a trademark of Myriad Women's Health, Inc.
Copyright 2018 Myriad Women's Health, Inc. All rights reserved.

INVITAE

INVITAE CARRIER SCREEN RESULTS

Patient name:	Donor 10289	**Sample type:**	Saliva	**Report date:**	27-JUN-2023
DOB:	[redacted]	**Sample collection date:**	19-JUN-2023	**Invitae #:**	RQ5202851
Sex assigned at birth:	Male	**Sample accession date:**	20-JUN-2023	**Clinical team:**	Guadalupe Martinez Dr. James Kuan
Gender:					
Patient ID (MRN):					

Reason for testing

Gamete donor

Test performed

Invitae Carrier Screen

RESULT: NEGATIVE

This carrier test evaluated 1 gene(s) for genetic changes (variants) that are associated with an increased risk of having a child with a genetic condition. Knowledge of carrier status for one of these conditions may provide information that can be used to assist with family planning and/or preparation. Carrier screening is not intended for diagnostic purposes. To identify a potential genetic basis for a condition in the individual being tested, diagnostic testing for the gene(s) of interest is recommended.

This test did not identify any genetic changes in the gene(s) analyzed that are currently recognized as clinically significant. This negative result reduces, but does not eliminate, the chance that this individual is a carrier for conditions caused by any of the genes tested. This individual may still be a carrier for a genetic condition that is not evaluated by this test.

Next steps

- Even for genes that have a negative test result, there is always a small risk that an individual could still be a carrier. This is called “residual risk.” See the Carrier detection rates and residual risks document.
- Discussion with a physician and/or genetic counselor is recommended to further review the implications of this test result and to understand these results in the context of any family history of a genetic condition.
- All patients, regardless of result, may wish to consider additional screening for hemoglobinopathies by complete blood count (CBC) and hemoglobin electrophoresis, if this has not already been completed.
- Individuals can register their tests at **https://www.invitae.com/patients/** to access online results, educational resources, and next steps.

Laboratory Director: Qing Zhang, MD, PhD, FACMG, NYCQ, CGMB
Invitae 1400 16th Street, San Francisco, CA 94103 · clientservices@invitae.com · 800.436.3037
© 2023 Invitae Corporation

Residual risk

No carrier test can detect 100% of carriers. There still remains a small risk of being a carrier after a negative test (residual risk). Residual risk values assume a negative family history and are inferred from published carrier frequencies and estimated detection rates based on testing technologies used at Invitae. You can view Invitae's complete Carrier detection rates and residual risks document (containing all carrier genes) online at **https://www.invitae.com/carrier-residual-risks/**. Additionally, the order-specific information for this report is available to download in the portal (under this order's documents) or can be requested by contacting Invitae Client Services. The complete Carrier detection rates and residual risks document will not be applicable for any genes with specimen-specific limitations in sequencing and/or deletion/duplication coverage. Please see the final bullet point in the Limitations section of this report to view if this specimen had any gene-specific coverage gaps.

© 2023 Invitae Corporation

Genes analyzed

This table represents a complete list of genes analyzed for this individual, including the relevant gene transcript(s). If more than one transcript is listed for a single gene, variants were reported using the first transcript listed unless otherwise indicated in the report. An asterisk (*) indicates that this gene has a limitation. Please see the Limitations section for details. Results are negative, unless otherwise indicated in the report.

GENE	TRANSCRIPT
DNAH5	NM_001369.2

© 2023 Invitae Corporation

Methods

- Genomic DNA obtained from the submitted sample is enriched for targeted regions using a hybridization-based protocol, and sequenced using Illumina technology. Unless otherwise indicated, all targeted regions are sequenced with ≥50x depth or are supplemented with additional analysis. Reads are aligned to a reference sequence (GRCh37), and sequence changes are identified and interpreted in the context of a single clinically relevant transcript, indicated in the Genes Analyzed table. Enrichment and analysis focus on the coding sequence of the indicated transcripts, 20bp of flanking intronic sequence, and other specific genomic regions demonstrated to be causative of disease at the time of assay design. Promoters, untranslated regions, and other non-coding regions are not otherwise interrogated. Exonic deletions and duplications are called using an in-house algorithm that determines copy number at each target by comparing the read depth for each target in the proband sequence with both mean read-depth and read-depth distribution, obtained from a set of clinical samples. Markers across the X and Y chromosomes are analyzed for quality control purposes and may detect deviations from the expected sex chromosome complement. Such deviations may be included in the report in accordance with internal guidelines. Invitae utilizes a classification methodology to identify next-generation sequencing (NGS)-detected variants that require orthogonal confirmation (Lincoln, et al. J Mol Diagn. 2019 Mar;21(2):318-329). Confirmation of the presence and location of reportable variants is performed as needed based on stringent criteria using one of several validated orthogonal approaches (PubMed ID 30610921). Sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778). Confirmatory sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778).

  The following additional analyses are performed if relevant to the requisition. For GBA the reference genome has been modified to mask the sites of polymorphic paralog sequence variants (PSVs) in both the gene and pseudogene. For CYP21A2 and GBA, if one or more reportable variants, gene conversion, or fusion event is identified via our NGS pipeline (see Limitations), these variants are confirmed by PacBio sequencing of an amplicon generated by long-range PCR and subsequent short-range PCR. In some cases, it may not be possible to disambiguate between the gene and pseudogene. For GJB2, the reportable range includes large upstream deletions overlapping GJB6. For HBA1/2, the reference genome has been modified to force some sequencing reads derived from HBA1 to align to HBA2, and variant calling algorithms are modified to support an expectation of 4 alleles in these regions. HBA1/2 copy number calling is performed by a custom hypothesis testing algorithm which generates diplotype calls. If sequence data for a sample does not support a unique high confidence match from among hypotheses tested, that sample is flagged for manual review. Copy number variation is only reported for coding sequence of HBA1 and HBA2 and the HS-40 region. This assay does not distinguish among the -α3.7 subtypes, and all -α3.7 variants are called as HBA1 deletions. This assay may not detect overlapping copy gain and copy loss events when the breakpoints of those events are similar. For FMR1, cytosine-guanine-guanine (CGG) triplet repeats in the 5' untranslated region (5' UTR) of the FMR1 gene are detected by triplet repeat-primed PCR (RP-PCR) with fluorescently labeled primers followed by capillary electrophoresis. Reference ranges: Normal: <45 CGG repeats, intermediate: 45-54 CGG repeats, premutation: 55-200 CGG repeats, full mutation: >200 CGG repeats. For alleles with 55-90 triplet repeats, the region surrounding the FMR1 repeat is amplified by PCR. The PCR amplicons are then processed through PacBio SMRTBell library prep and sequenced using PacBio long read technology. The number of AGG interruptions within the 55-90 triplet repeat is read directly from the resulting DNA sequences.

- This report only includes variants that have a clinically significant association with the conditions tested as of the report date. Variants of uncertain significance, benign variants, and likely benign variants are not included in this report. However, if additional evidence becomes available to indicate that the clinical significance of a variant has changed, Invitae may update this report and provide notification.
- A PMID is a unique identifier referring to a published, scientific paper. Search by PMID at **http://www.ncbi.nlm.nih.gov/pubmed**.
- An rsID is a unique identifier referring to a single genomic position, and is used to associate population frequency information with sequence changes at that position. Reported population frequencies are derived from a number of public sites that aggregate data from large-scale population sequencing projects, including ExAC (**http://exac.broadinstitute.org**), gnomAD (**http://gnomad.broadinstitute.org**), and dbSNP (**http://ncbi.nlm.nih.gov/SNP**).

Disclaimer

DNA studies do not constitute a definitive test for the selected condition(s) in all individuals. It should be realized that there are possible sources of error. Errors can result from trace contamination, rare technical errors, rare genetic variants that interfere with analysis, recent scientific developments, and alternative classification systems. This test should be one of many aspects used by the healthcare provider to help with a diagnosis and treatment plan, but it is not a diagnosis itself. This test was developed and its performance characteristics determined by Invitae. It has not been cleared or approved by

© 2023 Invitae Corporation

the FDA. The laboratory is regulated under the Clinical Laboratory Improvement Act (CLIA) as qualified to perform high-complexity clinical tests (CLIA ID: 05D2040778). This test is used for clinical purposes. It should not be regarded as investigational or for research.

Limitations

- Based on validation study results, this assay achieves >99% analytical sensitivity and specificity for single nucleotide variants, insertions and deletions <15bp in length, and exon-level deletions and duplications. Invitae's methods also detect insertions and deletions larger than 15bp but smaller than a full exon but sensitivity for these may be marginally reduced. Invitae's deletion/duplication analysis determines copy number at a single exon resolution at virtually all targeted exons. However, in rare situations, single-exon copy number events may not be analyzed due to inherent sequence properties or isolated reduction in data quality. Certain types of variants, such as structural rearrangements (e.g. inversions, gene conversion events, translocations, etc.) or variants embedded in sequence with complex architecture (e.g. short tandem repeats or segmental duplications), may not be detected. Additionally, it may not be possible to fully resolve certain details about variants, such as mosaicism, phasing, or mapping ambiguity. Unless explicitly guaranteed, sequence changes in the promoter, non-coding exons, and other non-coding regions are not covered by this assay. Please consult the test definition on our website for details regarding regions or types of variants that are covered or excluded for this test. This report reflects the analysis of an extracted genomic DNA sample. While this test is intended to reflect the analysis of extracted genomic DNA from a referred patient, in very rare cases the analyzed DNA may not represent that individual's constitutional genome, such as in the case of a circulating hematolymphoid neoplasm, bone marrow transplant, blood transfusion, chimerism, culture artifact or maternal cell contamination.

This report has been released utilizing a validated procedure approved by:

Qing Zhang, MD, PhD, FACMG, NYCQ, CGMB
Clinical Molecular Geneticist

© 2023 Invitae Corporation

INVITAE

CARRIER DETECTION RATES AND RESIDUAL RISKS

This table is relevant to patient report RQ5202851
Issue date: 06/27/2023

This table displays residual risks after a negative result for each of the genes and corresponding disorders. The values provided assume a negative family history and the absence of symptoms for each disorder. For genes associated with both dominant and recessive inheritance, the numbers in this table apply to the recessive condition(s) associated with the gene, unless otherwise noted. Residual risk values are provided for disorders when carrier frequency is greater than 1 in 500. For disorders with carrier frequency equal to, or less than, 1 in 500, residual risk is considered to be reduced substantially. When provided, residual risk values are inferred from published carrier frequencies, and estimated detection rates are based on testing technologies used at Invitae. Residual risks are provided only as a guide for assessing approximate risk given a negative result; values may vary based on the ethnic background(s) of an individual. For any genes marked with an asterisk*, refer to the Limitations section of the patient report for detailed coverage information. In the case of a sample-specific limitation, "N/A" indicates that a residual risk value could not be calculated. AR = autosomal recessive, XL = X-linked, AD = autosomal dominant.

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Primary ciliary dyskinesia (DNAH5-related) (AR) NM_001369.2	DNAH5	Pan-ethnic	1 in 109	99%	1 in 10800

WITHIN THE US | p: 800-436-3037 | f: 415-276-4164 | e: clientservices@invitae.com | OUTSIDE THE US | p: +1 415-930-4018 | f: +1 510 417-3922 | e: globalsupport@invitae.com
INVITAE | 1400 16th Street, San Francisco, CA 94103, USA | www.invitae.com/contact | © 2023 Invitae Corporation. All Rights Reserved.

Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
Phoenix Sperm Bank

1492 S Mill Ave Suite 306
Tempe, AZ 85281
Ph: (602)888-7255
AZB-45

LCLS Specimen Number:	**243-944-4014-0**	Account Number:	02003540
Patient Name:	**10289, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	08/31/2018
Lab Number:	YU18-70122 L	Date Received:	09/02/2018
Indications:	NOT GIVEN	Date Reported:	**09/19/2018**

Test: **Chromosome, Blood, Routine**

Cells Counted:	20	Cells Karyotyped:	2
Cells Analyzed:	20	Band Resolution:	500

CYTOGENETIC RESULT: 46,XY

INTERPRETATION: NORMAL MALE KARYOTYPE

Cytogenetic analysis of PHA stimulated cultures has revealed a MALE karyotype with an apparently normal GTG banding pattern in all cells observed.

This result does not exclude the possibility of subtle rearrangements below the resolution of cytogenetics or congenital anomalies due to other etiologies.

Chromosome analysis performed by LabCorp, CLIA 45D0674994. 7207 North Gessner Rd., Houston, TX 77040. Laboratory Director, Venkateswara R Potluri PhD.

Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
Phoenix Sperm Bank

1492 S Mill Ave Suite 306
Tempe, AZ 85281
Ph: (602)888-7255
AZB-45

LCLS Specimen Number:	**243-944-4014-0**
Patient Name:	**10289, DONOR**
Date of Birth:	[redacted]
Gender:	M
Patient ID:	
Lab Number:	YU18-70122 L

Account Number:	02003540
Ordering Physician:	**J OLLIFFE**
Specimen Type:	**BLOOD**
Client Reference:	
Date Collected:	08/31/2018
Date Received:	09/02/2018

Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
Phoenix Sperm Bank

1492 S Mill Ave Suite 306
Tempe, AZ 85281
Ph: (602)888-7255
AZB-45

LCLS Specimen Number:	**243-944-4014-0**	Account Number:	02003540
Patient Name:	**10289, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	08/31/2018
Lab Number:	YU18-70122 L	Date Received:	09/02/2018

John E. Wiley, Ph.D., FACMG.

Arundhati Chatterjee, MD
Medical Director
Stuart Schwartz, PhD
National Director of Cytogenetics

Technical component performed by Laboratory Corporation of America Holdings, 1904 TW Alexander Drive , RTP , NC , 27709-0153 (800) 345-4363

Professional Component performed by LabCorp CLIA 34D0655223, 2253 Stantonsburg Rd, Greenville, NC 27834. Laboratory Director, John E. Wiley, Ph.D.,FACMG
Integrated Genetics is a brand used by Esoterix Genetic Laboratories, LLC, a wholly-owned subsidiary of Laboratory Corporation of America Holdings.

This document contains private and confidential health information **protected by state and federal law.**

Patient: 10289, DONOR
DOB: [redacted]

Patient ID:

Control ID: B0080043174

Specimen ID: 243-944-4010-0
Date collected: 08/31/2018 1015 Local

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
			1/2 Avg.Risk	1.0 1.5	
			Avg.Risk	3.6 3.2	
			2X Avg.Risk	6.2 5.0	
			3X Avg.Risk	8.0 6.1	
Hemoglobin Fractionation					01
Hgb F	0.0		%	0.0 - 2.0	01
Hgb A	97.7		%	96.4 - 98.8	02
Hgb S	0.0		%	0.0	02
Hgb C	0.0		%	0.0	02
Hgb A2	2.3		%	1.8 - 3.2	02
Hgb Variant	0.0		%	0.0	02
Interpretation					02
Normal adult hemoglobin present.					
Serology/Immunology					01
ABO Grouping	B				01
Rh Factor	Positive				02
Please note: Prior records for this patient's ABO / Rh type are not available for additional verification.					01
CBC, Platelet Ct, and Diff					01
WBC	4.9		x10E3/uL	3.4 - 10.8	01
RBC	4.96		x10E6/uL	4.14 - 5.80	01
Hemoglobin	14.9		g/dL	13.0 - 17.7	01
Hematocrit	43.5		%	37.5 - 51.0	01
MCV	88		fL	79 - 97	01
MCH	30.0		pg	26.6 - 33.0	01
MCHC	34.3		g/dL	31.5 - 35.7	01
RDW	12.9		%	12.3 - 15.4	01
Platelets	278		x10E3/uL	150 - 379	01
Neutrophils	57		%	Not Estab.	01
Lymphs	34		%	Not Estab.	01
Monocytes	7		%	Not Estab.	01
Eos	2		%	Not Estab.	01
Basos	0		%	Not Estab.	01
Neutrophils (Absolute)	2.8		x10E3/uL	1.4 - 7.0	01
Lymphs (Absolute)	1.7		x10E3/uL	0.7 - 3.1	01
Monocytes(Absolute)	0.3		x10E3/uL	0.1 - 0.9	01
Eos (Absolute)	0.1		x10E3/uL	0.0 - 0.4	01
Baso (Absolute)	0.0		x10E3/uL	0.0 - 0.2	01
Immature Granulocytes	0		%	Not Estab.	01
Immature Grans (Abs)	0.0		x10E3/uL	0.0 - 0.1	01

Date Issued: 09/05/18 1642 ET

FINAL REPORT

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error, please call 800-788-9743

© 1995-2018 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version: 1.00