RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Jeffrey Olliffe
4915 25th Ave NE Ste 204w
Seattle, WA 98105-5668
Phone: (206) 588-1484
Fax: (206) 466-4696
NPI: 1306838271
Report Date: 03/05/2020

MALE
DONOR 12552
DOB: [redacted]
Ethnicity: Mixed or Other Caucasian
Sample Type: EDTA Blood
Date of Collection: 02/25/2020
Date Received: 02/27/2020
Date Tested: 03/03/2020
Barcode: 11004512621415
Accession ID: CSLXU3NEUEWJXPU
Indication: Egg or sperm donor

FEMALE
N/A

Foresight® Carrier Screen

POSITIVE: CARRIER

ABOUT THIS TEST

The **Myriad Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

RESULTS SUMMARY

Risk Details	DONOR 12552	Partner
Panel Information	Foresight Carrier Screen Universal Panel Fundamental Plus Panel Fundamental Panel **(175 conditions tested)**	N/A
POSITIVE: CARRIER Alpha Thalassemia **Reproductive Risk: Not Calculated** **Inheritance: Autosomal Recessive**	**CARRIER*** NM_000517.4(HBA2):c.427T>C (*143Qext*31, aka Hb Constant Spring) heterozygote Alpha globin status: -a/aa.	Reproductive risk can be more accurately assessed after carrier screening of the partner. Carrier testing should be considered. See "Next Steps".

*Carriers generally do not experience symptoms.

No disease-causing mutations were detected in any other gene tested. A complete list of all conditions tested can be found on page 7.

CLINICAL NOTES

- None

NEXT STEPS

- Carrier testing should be considered for the diseases specified above for the patient's partner, as both parents must be carriers before a child is at high risk of developing the disease.
- Genetic counseling is recommended and patients may wish to discuss any positive results with blood relatives, as there is an increased chance that they are also carriers.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

POSITIVE: CARRIER

Alpha Thalassemia

Genes: HBA1, HBA2 | **Inheritance Pattern:** Autosomal Recessive

	Patient DONOR 12552	**No partner tested**
Result	✚ Carrier	N/A
Variant(s)	NM_000517.4(HBA2):c.427T>C(*143Qext*31, aka Hb Constant Spring) heterozygote	N/A
Methodology	Analysis of homologous regions	N/A
Interpretation	This individual is a carrier of alpha thalassemia. Carriers do not experience symptoms, but may have hematologic abnormalities. Hb Constant Spring is classified as an alpha+ mutation. Based on this result, the patient's alpha globin status is -a/aa (carrier), where "-" indicates a deleted or nonfunctional alpha globin gene.	N/A
Detection rate	Unknown due to rarity of disease	N/A
Variants tested	-(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, anti3.7, anti4.2, del HS-40.	N/A

REPRODUCTIVE RISK SUMMARY
Reproductive risk can be more accurately assessed after carrier screening of the partner. Genetic counseling is recommended to review results and risks in further detail.

What is Alpha Thalassemia?

Alpha thalassemia is a blood disorder that affects hemoglobin, a major component of red blood cells that carries oxygen in the body. Hemoglobin is a protein complex made up of two different chains. There are many forms of hemoglobin, but the primary type is made up of alpha chains and beta chains. Alpha thalassemia is caused by mutations involving the genes, *HBA1* and *HBA2*, that code for the alpha chains.

Most individuals have two functional pairs or four functional copies of the alpha globin genes (one copy each of *HBA1* and *HBA2* on both chromosomes).

Carriers generally have either two or three functional alpha globin genes and do not have any symptoms.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

- **Three functional alpha globin genes, silent carrier:** These individuals are typically known as silent carriers, because they do not have any symptoms or abnormalities on a complete blood count. This status results from the presence of an alpha+ mutation (mutation that eliminates the function/presence of one copy of an alpha globin gene).

- **Two functional alpha globin genes, carrier:** These carriers generally have mild anemia characterized by hypochromic (pale) and microcytic (small) red blood cells, which can be measured on a complete blood count. However, they usually do not have any symptoms of the disease (note exception below). Carrier status may result from the presence of two alpha+ mutations (eliminates function/presence of one copy of an alpha globin gene on each chromosome) or an alpha0 mutation (eliminates function/presence of both copies of the alpha globin genes on one chromosome).

Exception: There have been reports of individuals with two copies of certain types of point mutations who have a diagnosis of hemoglobin H disease with variable symptoms. One example of this is when individuals have two copies of the hemoglobin Constant Spring mutation, which is common in the Southeast Asian population.

Disease symptoms most typically occur if an individual has one or zero functional alpha globin genes.

- **One functional alpha globin gene, hemoglobin H disease:** This form of alpha thalassemia is very variable. Disease severity ranges from asymptomatic to moderate microcytic/hypochromic anemia with the possibility of jaundice (yellowing of the skin or eyes), enlarged spleen, bone deformities, fatigue, and other minor complications.

- **Zero functional alpha globin genes, hemoglobin Bart syndrome:** Individuals who have no functional copies or are missing all four copies of the associated genes almost always have this fatal form of alpha thalassemia. Hb Bart syndrome is generally associated with death *in utero* due to the buildup of excess fluid in the body and tissues (hydrops fetalis). Signs and symptoms in the newborn period can include severe anemia, hepatosplenomegaly (enlarged liver and spleen), and birth defects of the heart, urinary system, and genitalia. Most babies with this condition are stillborn or die soon after birth.

How common is Alpha Thalassemia?

The carrier frequency and incidence of alpha thalassemia vary by the type and population. Carrier frequency of this condition is reported to be the highest in individuals of Southeast Asian, African, West Indian, and Mediterranean descent. In 2010, the estimated number of worldwide annual births of patients with Hb H disease was 9,568 and with Hb Bart syndrome was 5,183. Therefore, the worldwide birth prevalence of Hb H disease and Hb Bart's hydrops is estimated at ~1/14500 and ~1/27000, respectively; however, for Hb Bart's hydrops, this is likely to be an underestimate because most at-risk couples are not currently identified.

How is Alpha Thalassemia treated?

Alpha thalassemia carrier status does not necessitate treatment. Treatment for hemoglobin H disease varies based on the severity of the symptoms. For many individuals, blood transfusions are given during crises, which are episodic and usually precipitated by environmental stressors, like oxidant medications or fever. Individuals with more severe symptoms may require regular blood transfusions, folic acid supplementation, prophylactic antibiotics, iron chelation therapy (removal of excess iron from the body), and possible hemoglobin F-enhancing agents and splenectomy.

Extremely rare cases of survivors with hemoglobin Bart syndrome have been reported when fetal blood transfusions were given, followed by regular treatments similar to those who have hemoglobin H disease. Treatments or surgical correction of potential birth defects may also be available. However, there is a high risk for intellectual and physical disability in these rare survivors. These individuals may be candidates for hematopoietic stem cell transplantation.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

What is the prognosis for a person with Alpha Thalassemia?

Because hemoglobin H disease can be variable, prognosis ultimately depends on the severity of the disease. Mild disease may be manageable with little effect on daily life. However, more severe disease will necessitate frequent and regular therapy, and may be associated with a shortened lifespan. Untreated, the prognosis is poor with a shortened lifespan of up to age 5 years. However, when treated, individuals with hemoglobin H disease have a lifespan that approaches normal.

Hemoglobin Bart syndrome is the most severe clinical condition related to alpha thalassemia, and death may occur *in utero* or in the newborn period. Of note, there may also be maternal complications during pregnancy if the fetus has hemoglobin Bart syndrome. These complications include preeclampsia (high blood pressure, fluid build-up/swelling, protein in the urine), polyhydramnios (excessive amniotic fluid) or oligohydramnios (reduced amniotic fluid), hemorrhage, and premature delivery.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Methods and Limitations

DONOR 12552 [Foresight Carrier Screen]: Sequencing with copy number analysis, spinal muscular atrophy, and analysis of homologous regions (DTS v3).

Sequencing with copy number analysis

High-throughput sequencing and read depth-based copy number analysis are used to analyze the listed exons, as well as selected intergenic and intronic regions, of the genes in the Conditions Tested section of the report. The region of interest (ROI) of the test comprises these regions, in addition to the 20 intronic bases flanking each exon. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to high coverage and the sequences are compared to standards and references of normal variation (Genome Reference Consortium Human Build 37 (GRCh37)/hg19). More than 99% of all bases in the ROI are sequenced at greater than the minimum read depth. Mutations may not be detected in areas of lower sequence coverage. Small insertions and deletions may not be as accurately determined as single nucleotide variants. Genes that have closely related pseudogenes may be addressed by a different method. *CFTR* and *DMD* testing includes analysis for both large (exon-level) deletions and duplications with an average sensitivity of 99%, while other genes are only analyzed for large deletions with a sensitivity of >75%. However, the sensitivity may be higher for selected founder deletions. The breakpoints of copy number variants and exons affected are estimated from probe positions. Only exons known to be included in the copy number variant are provided in the name. In some cases, the copy number variant may be larger or smaller than indicated. If *GJB2* is tested, two large upstream deletions which overlap *GJB6* and affect the expression of *GJB2*, del(*GJB6*-D13S1830) and del(*GJB6*-D13S1854), are also analyzed. Mosaicism or somatic variants present at low levels may not be detected. If detected, these may not be reported.

Detection rates are determined by using literature to estimate the fraction of disease alleles, weighted by frequency, that the methodology is unable to detect. Detection rates only account for analytical sensitivity and certain variants that have been previously described in the literature may not be reported if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease-specific rates of de novo mutations.

All variants that are a recognized cause of the disease will be reported. In addition, variants that have not previously been established as a recognized cause of disease may be identified. In these cases, only variants classified as "likely" pathogenic are reported. Likely pathogenic variants are described elsewhere in the report as "likely to have a negative impact on gene function". Likely pathogenic variants are evaluated and classified by assessing the nature of the variant and reviewing reports of allele frequencies in cases and controls, functional studies, variant annotation and effect prediction, and segregation studies. Exon level duplications are assumed to be in tandem and are classified according to their predicted effect on the reading frame. Benign variants, variants of uncertain significance, and variants not directly associated with the intended disease phenotype are not reported. Curation summaries of reported variants are available upon request.

Spinal muscular atrophy

Targeted copy number analysis is used to determine the copy number of exon 7 of the *SMN1* gene relative to other genes. Other mutations may interfere with this analysis. Some individuals with two copies of *SMN1* are carriers with two *SMN1* genes on one chromosome and a *SMN1* deletion on the other chromosome. This is more likely in individuals who have 2 copies of the *SMN1* gene and are positive for the g.27134T>G SNP, which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have 2 copies of *SMN1*.

Analysis of homologous regions

A combination of high-throughput sequencing, read depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss of function mutations in certain genes that have homology to other regions. The precise breakpoints of large deletions in these genes cannot be determined, but are estimated from copy number analysis. High numbers of pseudogene copies may interfere with this analysis.

If *CYP21A2* is tested, patients who have one or more additional copies of the *CYP21A2* gene and a loss of function mutation may not actually be a carrier of 21-hydroxylase-deficient congenital adrenal hyperplasia (CAH). Because the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are only based on published incidences for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate of overall chances for 21-hydroxylase-deficient CAH, especially in the aforementioned populations, as they do not account for non-classic CAH. If *HBA1*/*HBA2* are tested, some individuals with four alpha globin genes may be carriers, with three genes on one chromosome and a deletion on the other chromosome. This and similar, but rare, carrier states, where complementary changes exist in both the gene and a pseudogene, may not be detected by the assay.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Limitations

In an unknown number of cases, nearby genetic variants may interfere with mutation detection. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions and technical errors. This test is designed to detect and report germline alterations. While somatic variants present at low levels may be detected, these may not be reported. f more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes. The test does not fully address all inherited forms of intellectual disability, birth defects and genetic disease. A family history of any of these conditions may warrant additional evaluation. Furthermore, not all mutations will be identified in the genes analyzed and additional testing may be beneficial for some patients. For example, individuals of African, Southeast Asian, and Mediterranean ancestry are at increased risk for being carriers for hemoglobinopathies, which can be identified by CBC and hemoglobin electrophoresis or HPLC (*ACOG Practice Bulletin No. 78. Obstet.Gynecol. 2007;109:229-37*).

This test was developed and its performance characteristics determined by Myriad Women's Health, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: **#05D1102604**.

Resources

GENOME CONNECT | http://www.genomeconnect.org

Patients can share their reports via research registries such as Genome Connect, an online research registry working to build the knowledge base about genetics and health. Genome Connect provides patients, physicians, and researchers an opportunity to share genetic information to support the study of the impact of genetic variation on health conditions.

SENIOR LABORATORY DIRECTOR

Jack Ji, PhD, FACMG

Report content approved by Jack Ji, PhD, FACMG on Mar 5, 2020

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Conditions Tested

11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia - **Gene:** CYP11B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000497:1-9. **Detection Rate:** Mixed or Other Caucasian 94%.

6-pyruvoyl-tetrahydropterin Synthase Deficiency - **Gene:** PTS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000317:1-6. **Detection Rate:** Mixed or Other Caucasian >99%.

ABCC8-related Familial Hyperinsulinism - **Gene:** ABCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000352:1-39. **Detection Rate:** Mixed or Other Caucasian >99%.

Adenosine Deaminase Deficiency - **Gene:** ADA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000022:1-12. **Detection Rate:** Mixed or Other Caucasian >99%.

Alpha Thalassemia - **Genes:** HBA1, HBA2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** -(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, anti3.7, anti4.2, del HS-40. **Detection Rate:** Unknown due to rarity of disease.

Alpha-mannosidosis - **Gene:** MAN2B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000528:1-23. **Detection Rate:** Mixed or Other Caucasian >99%.

Alpha-sarcoglycanopathy - **Gene:** SGCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000023:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.

Alstrom Syndrome - **Gene:** ALMS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015120:1-23. **Detection Rate:** Mixed or Other Caucasian >99%.

AMT-related Glycine Encephalopathy - **Gene:** AMT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000481:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.

Andermann Syndrome - **Gene:** SLC12A6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133647:1-25. **Detection Rate:** Mixed or Other Caucasian >99%.

Argininemia - **Gene:** ARG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000045:1-8. **Detection Rate:** Mixed or Other Caucasian 97%.

Argininosuccinic Aciduria - **Gene:** ASL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001024943:1-16. **Detection Rate:** Mixed or Other Caucasian >99%.

Aspartylglucosaminuria - **Gene:** AGA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000027:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.

Ataxia with Vitamin E Deficiency - **Gene:** TTPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000370:1-5. **Detection Rate:** Mixed or Other Caucasian >99%.

Ataxia-telangiectasia - **Gene:** ATM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000051:2-63. **Detection Rate:** Mixed or Other Caucasian 98%.

ATP7A-related Disorders - **Gene:** ATP7A. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000052:2-23. **Detection Rate:** Mixed or Other Caucasian 96%.

Autoimmune Polyglandular Syndrome Type 1 - **Gene:** AIRE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000383:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.

Autosomal Recessive Osteopetrosis Type 1 - **Gene:** TCIRG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006019:2-20. **Detection Rate:** Mixed or Other Caucasian >99%.

Autosomal Recessive Polycystic Kidney Disease, PKHD1-related - **Gene:** PKHD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138694 2-67. **Detection Rate:** Mixed or Other Caucasian >99%.

Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay - **Gene:** SACS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014363 2-10. **Detection Rate:** Mixed or Other Caucasian 99%.

Bardet-Biedl Syndrome, BBS1-related - **Gene:** BBS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024649:1-17. **Detection Rate:** Mixed or Other Caucasian >99%.

Bardet-Biedl Syndrome, BBS10-related - **Gene:** BBS10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024685:1-2. **Detection Rate:** Mixed or Other Caucasian >99%.

Bardet-Biedl Syndrome, BBS12-related - **Gene:** BBS12. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_152618:2. **Detection Rate:** Mixed or Other Caucasian >99%.

Bardet-Biedl Syndrome, BBS2-related - **Gene:** BBS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_031885:1-17. **Detection Rate:** Mixed or Other Caucasian >99%.

BCS1L-related Disorders - **Gene:** BCS1L. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004328:3-9. **Detection Rate:** Mixed or Other Caucasian >99%.

Beta-sarcoglycanopathy - **Gene:** SGCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000232:1-6. **Detection Rate:** Mixed or Other Caucasian >99%.

Biotinidase Deficiency - **Gene:** BTD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000060:1-4. **Detection Rate:** Mixed or Other Caucasian >99%.

Bloom Syndrome - **Gene:** BLM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000057:2-22. **Detection Rate:** Mixed or Other Caucasian >99%.

Calpainopathy - **Gene:** CAPN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000070:1-24. **Detection Rate:** Mixed or Other Caucasian >99%.

Canavan Disease - **Gene:** ASPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000049:1-6. **Detection Rate:** Mixed or Other Caucasian 98%.

Carbamoylphosphate Synthetase I Deficiency - **Gene:** CPS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001875:1-38. **Detection Rate:** Mixed or Other Caucasian >99%.

Carnitine Palmitoyltransferase IA Deficiency - **Gene:** CPT1A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001876:2-19. **Detection Rate:** Mixed or Other Caucasian >99%.

Carnitine Palmitoyltransferase II Deficiency - **Gene:** CPT2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000098:1-5. **Detection Rate:** Mixed or Other Caucasian >99%.

Cartilage-hair Hypoplasia - **Gene:** RMRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NR_003051:1. **Detection Rate:** Mixed or Other Caucasian >99%.

Cerebrotendinous Xanthomatosis - **Gene:** CYP27A1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000784:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.

Citrullinemia Type 1 - **Gene:** ASS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000050:3-16. **Detection Rate:** Mixed or Other Caucasian >99%.

CLN3-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001042432 2-16. **Detection Rate:** Mixed or Other Caucasian >99%.

CLN5-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006493:1-4. **Detection Rate:** Mixed or Other Caucasian >99%.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

CLN6-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017882:1-7. **Detection Rate:** Mixed or Other Caucasian >99%.
CLN8-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_018941:2-3. **Detection Rate:** Mixed or Other Caucasian >99%.
Cohen Syndrome - **Gene:** VPS13B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017890:2-62. **Detection Rate:** Mixed or Other Caucasian 97%.
COL4A3-related Alport Syndrome - **Gene:** COL4A3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000091:1-52. **Detection Rate:** Mixed or Other Caucasian 97%.
COL4A4-related Alport Syndrome - **Gene:** COL4A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000092:2-48. **Detection Rate:** Mixed or Other Caucasian 98%.
Combined Pituitary Hormone Deficiency, PROP1-related - **Gene:** PROP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006261:1-3. **Detection Rate:** Mixed or Other Caucasian >99%.
Congenital Adrenal Hyperplasia, CYP21A2-related - **Gene:** CYP21A2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs*21, I173N, L308Ffs*6, P31L, Q319*, Q319*+CYP21A2dup, R357W, V281L, [(I237N;V238E;M240K)], c.293-13C>G. **Detection Rate:** Mixed or Other Caucasian 96%.
Congenital Disorder of Glycosylation Type Ia - **Gene:** PMM2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000303:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Congenital Disorder of Glycosylation Type Ic - **Gene:** ALG6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_013339:2-15. **Detection Rate:** Mixed or Other Caucasian >99%.
Congenital Disorder of Glycosylation, MPI-related - **Gene:** MPI. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002435:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Costeff Optic Atrophy Syndrome - **Gene:** OPA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_025136:1-2. **Detection Rate:** Mixed or Other Caucasian >99%.
Cystic Fibrosis - **Gene:** CFTR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. **Detection Rate:** Mixed or Other Caucasian >99%.
Cystinosis - **Gene:** CTNS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004937:3-12. **Detection Rate:** Mixed or Other Caucasian >99%.
D-bifunctional Protein Deficiency - **Gene:** HSD17B4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000414:1-24. **Detection Rate:** Mixed or Other Caucasian 98%.
Delta-sarcoglycanopathy - **Gene:** SGCD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000337:2-9. **Detection Rate:** Mixed or Other Caucasian 99%.
Dihydrolipoamide Dehydrogenase Deficiency - **Gene:** DLD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000108:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.
Dysferlinopathy - **Gene:** DYSF. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003494:1-55. **Detection Rate:** Mixed or Other Caucasian 98%.
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy) - **Gene:** DMD. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_004006:1-79. **Detection Rate:** Mixed or Other Caucasian >99%.
ERCC6-related Disorders - **Gene:** ERCC6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000124:2-21. **Detection Rate:** Mixed or Other Caucasian 99%.
ERCC8-related Disorders - **Gene:** ERCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000082:1-12. **Detection Rate:** Mixed or Other Caucasian 95%.
EVC-related Ellis-van Creveld Syndrome - **Gene:** EVC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153717:1-21. **Detection Rate:** Mixed or Other Caucasian 96%.
EVC2-related Ellis-van Creveld Syndrome - **Gene:** EVC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_147127:1-22. **Detection Rate:** Mixed or Other Caucasian >99%.
Fabry Disease - **Gene:** GLA. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000169:1-7. **Detection Rate:** Mixed or Other Caucasian 98%.
Familial Dysautonomia - **Gene:** IKBKAP. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003640:2-37. **Detection Rate:** Mixed or Other Caucasian >99%.
Familial Mediterranean Fever - **Gene:** MEFV. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000243:1-10. **Detection Rate:** Mixed or Other Caucasian >99%.
Fanconi Anemia Complementation Group A - **Gene:** FANCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000135:1-43. **Detection Rate:** Mixed or Other Caucasian 92%.
Fanconi Anemia, FANCC-related - **Gene:** FANCC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000136:2-15. **Detection Rate:** Mixed or Other Caucasian >99%.
FKRP-related Disorders - **Gene:** FKRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_024301:4. **Detection Rate:** Mixed or Other Caucasian >99%.
FKTN-related Disorders - **Gene:** FKTN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001079802:3-11. **Detection Rate:** Mixed or Other Caucasian >99%.
Galactokinase Deficiency - **Gene:** GALK1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000154:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Galactosemia - **Gene:** GALT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000155:1-11. **Detection Rate:** Mixed or Other Caucasian >99%.
Gamma-sarcoglycanopathy - **Gene:** SGCG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000231:2-8. **Detection Rate:** Mixed or Other Caucasian 88%.
Gaucher Disease - **Gene:** GBA. Autosomal Recessive. Analysis of homologous regions. **Variants (10):** D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs*18. **Detection Rate:** Mixed or Other Caucasian 60%.
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness - **Gene:** GJB2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004004:1-2. **Detection Rate:** Mixed or Other Caucasian >99%.
GLB1-related Disorders - **Gene:** GLB1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000404:1-16. **Detection Rate:** Mixed or Other Caucasian >99%.
GLDC-related Glycine Encephalopathy - **Gene:** GLDC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000170:1-25. **Detection Rate:** Mixed or Other Caucasian 94%.
Glutaric Acidemia, GCDH-related - **Gene:** GCDH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000159:2-12. **Detection Rate:** Mixed or Other Caucasian >99%.
Glycogen Storage Disease Type Ia - **Gene:** G6PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000151:1-5. **Detection Rate:** Mixed or Other Caucasian >99%.
Glycogen Storage Disease Type Ib - **Gene:** SLC37A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001164277 3-11. **Detection Rate:** Mixed or Other Caucasian >99%.
Glycogen Storage Disease Type III - **Gene:** AGL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000642:2-34. **Detection Rate:** Mixed or Other Caucasian >99%.
GNE Myopathy - **Gene:** GNE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001128227:1-12. **Detection Rate:** Mixed or Other Caucasian >99%.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

GNPTAB-related Disorders - **Gene:** GNPTAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024312:1-21. **Detection Rate:** Mixed or Other Caucasian >99%.
HADHA-related Disorders - **Gene:** HADHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000182:1-20. **Detection Rate:** Mixed or Other Caucasian >99%.
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease) - **Gene:** HBB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000518:1-3. **Detection Rate:** Mixed or Other Caucasian >99%.
Hereditary Fructose Intolerance - **Gene:** ALDOB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000035:2-9. **Detection Rate:** Mixed or Other Caucasian >99%.
Herlitz Junctional Epidermolysis Bullosa, LAMB3-related - **Gene:** LAMB3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000228 2-23. **Detection Rate:** Mixed or Other Caucasian >99%.
Hexosaminidase A Deficiency (Including Tay-Sachs Disease) - **Gene:** HEXA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000520:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.
HMG-CoA Lyase Deficiency - **Gene:** HMGCL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000191:1-9. **Detection Rate:** Mixed or Other Caucasian 98%.
Holocarboxylase Synthetase Deficiency - **Gene:** HLCS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000411:4-12. **Detection Rate:** Mixed or Other Caucasian >99%.
Homocystinuria, CBS-related - **Gene:** CBS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000071:3-17. **Detection Rate:** Mixed or Other Caucasian >99%.
Hydrolethalus Syndrome - **Gene:** HYLS1. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_145014:4. **Detection Rate:** Mixed or Other Caucasian >99%.
Hypophosphatasia - **Gene:** ALPL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000478:2-12. **Detection Rate:** Mixed or Other Caucasian >99%.
Isovaleric Acidemia - **Gene:** IVD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002225:1-12. **Detection Rate:** Mixed or Other Caucasian >99%.
Joubert Syndrome 2 - **Gene:** TMEM216. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001173990:1-5. **Detection Rate:** Mixed or Other Caucasian >99%.
Junctional Epidermolysis Bullosa, LAMA3-related - **Gene:** LAMA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000227:1-38. **Detection Rate:** Mixed or Other Caucasian >99%.
Junctional Epidermolysis Bullosa, LAMC2-related - **Gene:** LAMC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005562:1-23. **Detection Rate:** Mixed or Other Caucasian >99%.
KCNJ11-related Familial Hyperinsulinism - **Gene:** KCNJ11. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000525:1. **Detection Rate:** Mixed or Other Caucasian >99%.
Krabbe Disease - **Gene:** GALC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000153:1-17. **Detection Rate:** Mixed or Other Caucasian >99%.
LAMA2-related Muscular Dystrophy - **Gene:** LAMA2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000426:1-65. **Detection Rate:** Mixed or Other Caucasian >99%.
Leigh Syndrome, French-Canadian Type - **Gene:** LRPPRC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133259:1-38. **Detection Rate:** Mixed or Other Caucasian >99%.
Lipoid Congenital Adrenal Hyperplasia - **Gene:** STAR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000349:1-7. **Detection Rate:** Mixed or Other Caucasian >99%.
Lysosomal Acid Lipase Deficiency - **Gene:** LIPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000235:2-10. **Detection Rate:** Mixed or Other Caucasian >99%.
Maple Syrup Urine Disease Type Ia - **Gene:** BCKDHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000709:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.
Maple Syrup Urine Disease Type Ib - **Gene:** BCKDHB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_183050:1-10. **Detection Rate:** Mixed or Other Caucasian >99%.
Maple Syrup Urine Disease Type II - **Gene:** DBT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001918:1-11. **Detection Rate:** Mixed or Other Caucasian 96%.
Medium Chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000016:1-12. **Detection Rate:** Mixed or Other Caucasian >99%.
Megalencephalic Leukoencephalopathy with Subcortical Cysts - **Gene:** MLC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015166 2-12. **Detection Rate:** Mixed or Other Caucasian >99%.
Metachromatic Leukodystrophy - **Gene:** ARSA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000487:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Methylmalonic Acidemia, cblA Type - **Gene:** MMAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_172250:2-7. **Detection Rate:** Mixed or Other Caucasian >99%.
Methylmalonic Acidemia, cblB Type - **Gene:** MMAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_052845:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.
Methylmalonic Aciduria and Homocystinuria, cblC Type - **Gene:** MMACHC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015506:1-4. **Detection Rate:** Mixed or Other Caucasian >99%.
MKS1-related Disorders - **Gene:** MKS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017777:1-18. **Detection Rate:** Mixed or Other Caucasian >99%.
Mucolipidosis III Gamma - **Gene:** GNPTG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032520:1-11. **Detection Rate:** Mixed or Other Caucasian >99%.
Mucolipidosis IV - **Gene:** MCOLN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_020533:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.
Mucopolysaccharidosis Type I - **Gene:** IDUA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000203:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.
Mucopolysaccharidosis Type II - **Gene:** IDS. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000202:1-9. **Detection Rate:** Mixed or Other Caucasian 88%.
Mucopolysaccharidosis Type IIIA - **Gene:** SGSH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000199:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Mucopolysaccharidosis Type IIIB - **Gene:** NAGLU. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000263:1-6. **Detection Rate:** Mixed or Other Caucasian >99%.
Mucopolysaccharidosis Type IIIC - **Gene:** HGSNAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_152419:1-18. **Detection Rate:** Mixed or Other Caucasian >99%.
MUT-related Methylmalonic Acidemia - **Gene:** MUT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000255:2-13. **Detection Rate:** Mixed or Other Caucasian >99%.
MYO7A-related Disorders - **Gene:** MYO7A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000260:2-49. **Detection Rate:** Mixed or Other Caucasian >99%.
NEB-related Nemaline Myopathy - **Gene:** NEB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001271208:3-80,117-183. **Detection Rate:** Mixed or Other Caucasian 92%.
Nephrotic Syndrome, NPHS1-related - **Gene:** NPHS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004646:1-29. **Detection Rate:** Mixed or Other Caucasian >99%.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Nephrotic Syndrome, NPHS2-related - **Gene:** NPHS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014625:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Niemann-Pick Disease Type C1 - **Gene:** NPC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000271:1-25. **Detection Rate:** Mixed or Other Caucasian >99%.
Niemann-Pick Disease Type C2 - **Gene:** NPC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006432:1-5. **Detection Rate:** Mixed or Other Caucasian >99%.
Niemann-Pick Disease, SMPD1-related - **Gene:** SMPD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000543:1-6. **Detection Rate:** Mixed or Other Caucasian >99%.
Nijmegen Breakage Syndrome - **Gene:** NBN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002485:1-16. **Detection Rate:** Mixed or Other Caucasian >99%.
Ornithine Transcarbamylase Deficiency - **Gene:** OTC. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000531:1-10. **Detection Rate:** Mixed or Other Caucasian 97%.
PCCA-related Propionic Acidemia - **Gene:** PCCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000282:1-24. **Detection Rate:** Mixed or Other Caucasian 95%.
PCCB-related Propionic Acidemia - **Gene:** PCCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000532:1-15. **Detection Rate:** Mixed or Other Caucasian >99%.
PCDH15-related Disorders - **Gene:** PCDH15. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_033056:2-33. **Detection Rate:** Mixed or Other Caucasian 93%.
Pendred Syndrome - **Gene:** SLC26A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000441:2-21. **Detection Rate:** Mixed or Other Caucasian >99%.
Peroxisome Biogenesis Disorder Type 1 - **Gene:** PEX1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000466:1-24. **Detection Rate:** Mixed or Other Caucasian >99%.
Peroxisome Biogenesis Disorder Type 3 - **Gene:** PEX12. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000286:1-3. **Detection Rate:** Mixed or Other Caucasian >99%.
Peroxisome Biogenesis Disorder Type 4 - **Gene:** PEX6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000287:1-17. **Detection Rate:** Mixed or Other Caucasian 97%.
Peroxisome Biogenesis Disorder Type 5 - **Gene:** PEX2. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000318:4. **Detection Rate:** Mixed or Other Caucasian >99%.
Peroxisome Biogenesis Disorder Type 6 - **Gene:** PEX10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153818:1-6. **Detection Rate:** Mixed or Other Caucasian >99%.
Phenylalanine Hydroxylase Deficiency - **Gene:** PAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000277:1-13. **Detection Rate:** Mixed or Other Caucasian >99%.
POMGNT-related Disorders - **Gene:** POMGNT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017739:2-22. **Detection Rate:** Mixed or Other Caucasian 96%.
Pompe Disease - **Gene:** GAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000152:2-20. **Detection Rate:** Mixed or Other Caucasian 98%.
PPT1-related Neuronal Ceroid Lipofuscinosis - **Gene:** PPT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000310:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.
Primary Carnitine Deficiency - **Gene:** SLC22A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003060:1-10. **Detection Rate:** Mixed or Other Caucasian >99%.
Primary Hyperoxaluria Type 1 - **Gene:** AGXT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000030:1-11. **Detection Rate:** Mixed or Other Caucasian >99%.
Primary Hyperoxaluria Type 2 - **Gene:** GRHPR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012203:1-9. **Detection Rate:** Mixed or Other Caucasian >99%.
Primary Hyperoxaluria Type 3 - **Gene:** HOGA1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138413:1-7. **Detection Rate:** Mixed or Other Caucasian >99%.
Pycnodysostosis - **Gene:** CTSK. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000396:2-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Pyruvate Carboxylase Deficiency - **Gene:** PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000920:3-22. **Detection Rate:** Mixed or Other Caucasian >99%.
Rhizomelic Chondrodysplasia Punctata Type 1 - **Gene:** PEX7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000288:1-10. **Detection Rate:** Mixed or Other Caucasian >99%.
RTEL1-related Disorders - **Gene:** RTEL1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032957:2-35. **Detection Rate:** Mixed or Other Caucasian >99%.
Salla Disease - **Gene:** SLC17A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012434:1-11. **Detection Rate:** Mixed or Other Caucasian 98%.
Sandhoff Disease - **Gene:** HEXB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000521:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.
Short-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000017:1-10. **Detection Rate:** Mixed or Other Caucasian >99%.
Sjogren-Larsson Syndrome - **Gene:** ALDH3A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000382:1-10. **Detection Rate:** Mixed or Other Caucasian 96%.
SLC26A2-related Disorders - **Gene:** SLC26A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000112:2-3. **Detection Rate:** Mixed or Other Caucasian >99%.
Smith-Lemli-Opitz Syndrome - **Gene:** DHCR7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001360:3-9. **Detection Rate:** Mixed or Other Caucasian >99%.
Spastic Paraplegia Type 15 - **Gene:** ZFYVE26. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015346:2-42. **Detection Rate:** Mixed or Other Caucasian >99%.
Spinal Muscular Atrophy - **Gene:** SMN1. Autosomal Recessive. Spinal muscular atrophy. **Variant (1):** SMN1 copy number. **Detection Rate:** Mixed or Other Caucasian 95%.
Spondylothoracic Dysostosis - **Gene:** MESP2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001039958:1-2. **Detection Rate:** Mixed or Other Caucasian >99%.
TGM1-related Autosomal Recessive Congenital Ichthyosis - **Gene:** TGM1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000359 2-15. **Detection Rate:** Mixed or Other Caucasian >99%.
TPP1-related Neuronal Ceroid Lipofuscinosis - **Gene:** TPP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000391:1-13. **Detection Rate:** Mixed or Other Caucasian >99%.
Tyrosine Hydroxylase Deficiency - **Gene:** TH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_199292:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.
Tyrosinemia Type I - **Gene:** FAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000137:1-14. **Detection Rate:** Mixed or Other Caucasian >99%.
Tyrosinemia Type II - **Gene:** TAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000353:2-12. **Detection Rate:** Mixed or Other Caucasian >99%.
USH1C-related Disorders - **Gene:** USH1C. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005709:1-21. **Detection Rate:** Mixed or Other Caucasian >99%.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

USH2A-related Disorders - **Gene:** USH2A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_206933:2-72. **Detection Rate:** Mixed or Other Caucasian 94%.
Usher Syndrome Type 3 - **Gene:** CLRN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_174878:1-3. **Detection Rate:** Mixed or Other Caucasian >99%.
Very-long-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADVL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000018:1-20. **Detection Rate:** Mixed or Other Caucasian >99%.
Wilson Disease - **Gene:** ATP7B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000053:1-21. **Detection Rate:** Mixed or Other Caucasian >99%.
X-linked Adrenoleukodystrophy - **Gene:** ABCD1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000033:1-6. **Detection Rate:** Mixed or Other Caucasian 77%.
X-linked Alport Syndrome - **Gene:** COL4A5. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000495:1-51. **Detection Rate:** Mixed or Other Caucasian 95%.
X-linked Congenital Adrenal Hypoplasia - **Gene:** NR0B1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000475:1-2. **Detection Rate:** Mixed or Other Caucasian 99%.
X-linked Juvenile Retinoschisis - **Gene:** RS1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000330:1-6. **Detection Rate:** Mixed or Other Caucasian 98%.
X-linked Myotubular Myopathy - **Gene:** MTM1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000252:2-15. **Detection Rate:** Mixed or Other Caucasian 98%.
X-linked Severe Combined Immunodeficiency - **Gene:** IL2RG. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000206:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.
Xeroderma Pigmentosum Group A - **Gene:** XPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000380:1-6. **Detection Rate:** Mixed or Other Caucasian >99%.
Xeroderma Pigmentosum Group C - **Gene:** XPC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004628:1-16. **Detection Rate:** Mixed or Other Caucasian 97%.

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Risk Calculations

Below are the risk calculations for all conditions tested. Since negative results do not completely rule out the possibility of being a carrier, the **residual risk** represents the patient's post-test likelihood of being a carrier and the **reproductive risk** represents the likelihood the patient's future children could inherit each disease. These risks are inherent to all carrier screening tests, may vary by ethnicity, are predicated on a negative family history and are present even after a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

†Indicates a positive result. See the full clinical report for interpretation and details.

Disease	DONOR 12552 Residual Risk	Reproductive Risk
11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia	1 in 3,800	< 1 in 1,000,000
6-pyruvoyl-tetrahydropterin Synthase Deficiency	< 1 in 50,000	< 1 in 1,000,000
ABCC8-related Familial Hyperinsulinism	1 in 17,000	< 1 in 1,000,000
Adenosine Deaminase Deficiency	1 in 22,000	< 1 in 1,000,000
Alpha Thalassemia	NM_000517.4(HBA2):c.427T>C(*143Qext*31, aka Hb Constant Spring) heterozygote † Alpha globin status: -a/aa.	Not calculated
Alpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Alpha-sarcoglycanopathy	1 in 45,000	< 1 in 1,000,000
Alstrom Syndrome	< 1 in 50,000	< 1 in 1,000,000
AMT-related Glycine Encephalopathy	1 in 22,000	< 1 in 1,000,000
Andermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
Argininemia	< 1 in 17,000	< 1 in 1,000,000
Argininosuccinic Aciduria	1 in 13,000	< 1 in 1,000,000
Aspartylglucosaminuria	< 1 in 50,000	< 1 in 1,000,000
Ataxia with Vitamin E Deficiency	< 1 in 50,000	< 1 in 1,000,000
Ataxia-telangiectasia	1 in 11,000	< 1 in 1,000,000
ATP7A-related Disorders	< 1 in 1,000,000	1 in 600,000
Autoimmune Polyglandular Syndrome Type 1	1 in 15,000	< 1 in 1,000,000
Autosomal Recessive Osteopetrosis Type 1	1 in 35,000	< 1 in 1,000,000
Autosomal Recessive Polycystic Kidney Disease, PKHD1-related	1 in 8,100	< 1 in 1,000,000
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	< 1 in 44,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS1-related	1 in 32,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS10-related	1 in 42,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS12-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS2-related	< 1 in 50,000	< 1 in 1,000,000
BCS1L-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Beta-sarcoglycanopathy	1 in 39,000	< 1 in 1,000,000
Biotinidase Deficiency	1 in 13,000	1 in 650,000
Bloom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Calpainopathy	1 in 13,000	< 1 in 1,000,000
Canavan Disease	1 in 9,700	< 1 in 1,000,000
Carbamoylphosphate Synthetase I Deficiency	< 1 in 57,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase II Deficiency	1 in 25,000	< 1 in 1,000,000
Cartilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
Cerebrotendinous Xanthomatosis	1 in 11,000	< 1 in 1,000,000
Citrullinemia Type 1	1 in 14,000	< 1 in 1,000,000
CLN3-related Neuronal Ceroid Lipofuscinosis	1 in 8,600	< 1 in 1,000,000
CLN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN6-related Neuronal Ceroid Lipofuscinosis	1 in 43,000	< 1 in 1,000,000
CLN8-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Cohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
COL4A3-related Alport Syndrome	1 in 6,200	< 1 in 1,000,000
COL4A4-related Alport Syndrome	1 in 12,000	< 1 in 1,000,000
Combined Pituitary Hormone Deficiency, PROP1-related	1 in 6,100	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP21A2-related	1 in 1,300	1 in 280,000
Congenital Disorder of Glycosylation Type Ia	1 in 16,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ic	< 1 in 50,000	< 1 in 1,000,000

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 12552 Residual Risk	Reproductive Risk
Congenital Disorder of Glycosylation, MPI-related	< 1 in 50,000	< 1 in 1,000,000
Costeff Optic Atrophy Syndrome	< 1 in 50,000	< 1 in 1,000,000
Cystic Fibrosis	1 in 3,000	1 in 360,000
Cystinosis	1 in 22,000	< 1 in 1,000,000
D-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000
Delta-sarcoglycanopathy	< 1 in 40,000	< 1 in 1,000,000
Dihydrolipoamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Dysferlinopathy	1 in 11,000	< 1 in 1,000,000
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)	Not calculated	Not calculated
ERCC6-related Disorders	1 in 26,000	< 1 in 1,000,000
ERCC8-related Disorders	< 1 in 9,900	< 1 in 1,000,000
EVC-related Ellis-van Creveld Syndrome	1 in 7,500	< 1 in 1,000,000
EVC2-related Ellis-van Creveld Syndrome	< 1 in 50,000	< 1 in 1,000,000
Fabry Disease	< 1 in 1,000,000	1 in 80,000
Familial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
Familial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
Fanconi Anemia Complementation Group A	1 in 2,800	< 1 in 1,000,000
Fanconi Anemia, FANCC-related	< 1 in 50,000	< 1 in 1,000,000
FKRP-related Disorders	1 in 16,000	< 1 in 1,000,000
FKTN-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Galactokinase Deficiency	1 in 10,000	< 1 in 1,000,000
Galactosemia	1 in 8,600	< 1 in 1,000,000
Gamma-sarcoglycanopathy	1 in 3,000	< 1 in 1,000,000
Gaucher Disease	1 in 260	1 in 110,000
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 2,500	1 in 260,000
GLB1-related Disorders	1 in 19,000	< 1 in 1,000,000
GLDC-related Glycine Encephalopathy	1 in 2,800	< 1 in 1,000,000
Glutaric Acidemia, GCDH-related	1 in 16,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ia	1 in 18,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ib	1 in 35,000	< 1 in 1,000,000
Glycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
GNE Myopathy	1 in 23,000	< 1 in 1,000,000
GNPTAB-related Disorders	1 in 32,000	< 1 in 1,000,000
HADHA-related Disorders	1 in 20,000	< 1 in 1,000,000
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease)	1 in 3,100	1 in 390,000
Hereditary Fructose Intolerance	1 in 7,900	< 1 in 1,000,000
Herlitz Junctional Epidermolysis Bullosa, LAMB3-related	< 1 in 50,000	< 1 in 1,000,000
Hexosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
HMG-CoA Lyase Deficiency	< 1 in 33,000	< 1 in 1,000,000
Holocarboxylase Synthetase Deficiency	1 in 15,000	< 1 in 1,000,000
Homocystinuria, CBS-related	1 in 9,400	< 1 in 1,000,000
Hydrolethalus Syndrome	< 1 in 50,000	< 1 in 1,000,000
Hypophosphatasia	1 in 27,000	< 1 in 1,000,000
Isovaleric Acidemia	1 in 32,000	< 1 in 1,000,000
Joubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
KCNJ11-related Familial Hyperinsulinism	< 1 in 50,000	< 1 in 1,000,000
Krabbe Disease	1 in 14,000	< 1 in 1,000,000
LAMA2-related Muscular Dystrophy	1 in 34,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type	< 1 in 50,000	< 1 in 1,000,000
Lipoid Congenital Adrenal Hyperplasia	< 1 in 50,000	< 1 in 1,000,000
Lysosomal Acid Lipase Deficiency	1 in 18,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ia	1 in 42,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ib	1 in 39,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type II	1 in 13,000	< 1 in 1,000,000
Medium Chain Acyl-CoA Dehydrogenase Deficiency	1 in 4,400	1 in 790,000
Megalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000	< 1 in 1,000,000
Metachromatic Leukodystrophy	1 in 16,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblA Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblB Type	1 in 48,000	< 1 in 1,000,000

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

myriad WOMEN'S HEALTH

RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Jeffrey Olliffe
NPI: 1306838271
Report Date: 03/05/2020

MALE
DONOR 12552
DOB: [redacted]
Ethnicity: Mixed or Other Caucasian
Barcode: 11004512621415

FEMALE
N/A

Disease	DONOR 12552 Residual Risk	Reproductive Risk
Methylmalonic Aciduria and Homocystinuria, cblC Type	1 in 16,000	< 1 in 1,000,000
MKS1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis III Gamma	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis IV	< 1 in 50,000	< 1 in 1,000,000
Mucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type II	1 in 600,000	1 in 150,000
Mucopolysaccharidosis Type IIIA	1 in 12,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIB	1 in 25,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIC	1 in 37,000	< 1 in 1,000,000
MUT-related Methylmalonic Acidemia	1 in 26,000	< 1 in 1,000,000
MYO7A-related Disorders	1 in 15,000	< 1 in 1,000,000
NEB-related Nemaline Myopathy	1 in 1,200	1 in 400,000
Nephrotic Syndrome, NPHS1-related	< 1 in 50,000	< 1 in 1,000,000
Nephrotic Syndrome, NPHS2-related	1 in 35,000	< 1 in 1,000,000
Niemann-Pick Disease Type C1	1 in 19,000	< 1 in 1,000,000
Niemann-Pick Disease Type C2	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease, SMPD1-related	1 in 25,000	< 1 in 1,000,000
Nijmegen Breakage Syndrome	1 in 16,000	< 1 in 1,000,000
Ornithine Transcarbamylase Deficiency	< 1 in 1,000,000	1 in 140,000
PCCA-related Propionic Acidemia	1 in 4,200	< 1 in 1,000,000
PCCB-related Propionic Acidemia	1 in 22,000	< 1 in 1,000,000
PCDH15-related Disorders	1 in 3,300	< 1 in 1,000,000
Pendred Syndrome	1 in 8,200	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 1	1 in 16,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 3	1 in 44,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 4	1 in 9,300	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 5	< 1 in 71,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 6	< 1 in 50,000	< 1 in 1,000,000
Phenylalanine Hydroxylase Deficiency	1 in 4,800	1 in 940,000
POMGNT-related Disorders	< 1 in 12,000	< 1 in 1,000,000
Pompe Disease	1 in 4,000	< 1 in 1,000,000
PPT1-related Neuronal Ceroid Lipofuscinosis	1 in 7,700	< 1 in 1,000,000
Primary Carnitine Deficiency	1 in 11,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 1	1 in 17,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 3	1 in 13,000	< 1 in 1,000,000
Pycnodysostosis	1 in 43,000	< 1 in 1,000,000
Pyruvate Carboxylase Deficiency	1 in 25,000	< 1 in 1,000,000
Rhizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
RTEL1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Salla Disease	< 1 in 30,000	< 1 in 1,000,000
Sandhoff Disease	1 in 32,000	< 1 in 1,000,000
Short-chain Acyl-CoA Dehydrogenase Deficiency	1 in 11,000	< 1 in 1,000,000
Sjogren-Larsson Syndrome	< 1 in 12,000	< 1 in 1,000,000
SLC26A2-related Disorders	1 in 16,000	< 1 in 1,000,000
Smith-Lemli-Opitz Syndrome	1 in 9,400	< 1 in 1,000,000
Spastic Paraplegia Type 15	< 1 in 50,000	< 1 in 1,000,000
Spinal Muscular Atrophy	Negative for g.27134T>G SNP SMN1: 2 copies 1 in 770	1 in 110,000
Spondylothoracic Dysostosis	< 1 in 50,000	< 1 in 1,000,000
TGM1-related Autosomal Recessive Congenital Ichthyosis	1 in 22,000	< 1 in 1,000,000
TPP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
Tyrosine Hydroxylase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Tyrosinemia Type I	1 in 16,000	< 1 in 1,000,000
Tyrosinemia Type II	1 in 25,000	< 1 in 1,000,000
USH1C-related Disorders	1 in 35,000	< 1 in 1,000,000
USH2A-related Disorders	1 in 2,200	< 1 in 1,000,000
Usher Syndrome Type 3	1 in 41,000	< 1 in 1,000,000
Very-long-chain Acyl-CoA Dehydrogenase Deficiency	1 in 18,000	< 1 in 1,000,000
Wilson Disease	1 in 8,600	< 1 in 1,000,000
X-linked Adrenoleukodystrophy	1 in 90,000	1 in 42,000

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 12552 Residual Risk	Reproductive Risk
X-linked Alport Syndrome	Not calculated	Not calculated
X-linked Congenital Adrenal Hypoplasia	< 1 in 1,000,000	< 1 in 1,000,000
X-linked Juvenile Retinoschisis	< 1 in 1,000,000	1 in 40,000
X-linked Myotubular Myopathy	Not calculated	Not calculated
X-linked Severe Combined Immunodeficiency	< 1 in 1,000,000	1 in 200,000
Xeroderma Pigmentosum Group A	< 1 in 50,000	< 1 in 1,000,000
Xeroderma Pigmentosum Group C	1 in 7,300	< 1 in 1,000,000

Copyright 2020 Myriad Women's Health, Inc. All rights reserved.

Patient Information
Name: Donor 12552
Date of Birth [redacted]
Sema4 ID: 22216546CS
Client ID: SEATSB-S411956848
Indication: Carrier Screening

Specimen Information
Specimen Type: Blood
Date Collected: 10/25/2022
Date Received: 10/26/2022
Final Report: 11/08/2022

Referring Provider
Jeffrey Olliffe, M.D.
Seattle Sperm Bank
4915 25th Avenue NE Suite 204W
Seattle, WA, 98105
Fax: 206-466-4696

Custom Carrier Screen (1 gene)

with Personalized Residual Risk

SUMMARY OF RESULTS AND RECOMMENDATIONS

Negative

Negative for all genes tested: *ABCA12*
To view a full list of genes and diseases tested
please see Table 1 in this report

AR=Autosomal recessive; XL=X-linked

Recommendations

- Consideration of residual risk by ethnicity after a negative carrier screen is recommended for the other diseases on the panel, especially in the case of a positive family history for a specific disorder. Please note that residual risks for X-linked diseases (including full repeat expansions for Fragile X syndrome) may not be accurate for males and the actual residual risk is likely to be lower.
- As genetic technologies may improve and variant classifications may change over time, it is recommended to obtain a new carrier screening test or reanalysis when a new pregnancy is being considered.

Test description

This patient was tested for the genes listed above using one or more of the following methodologies: target capture and short-read sequencing, long-range PCR followed by short-read sequencing, targeted genotyping, and/or copy number analysis. Please note that negative results reduce but do not eliminate the possibility that this individual is a carrier for one or more of the disorders tested. Please see Table 1 for a list of genes and diseases tested with the patient's personalized residual risk. If personalized residual risk is not provided, please see the complete residual risk table at **go.sema4.com/residualrisk.** Only known pathogenic or likely pathogenic variants are reported. This carrier screening test does not report likely benign variants and variants of uncertain significance (VUS). If reporting of likely benign variants and VUS are desired in this patient, please contact the laboratory at 800-298-6470, option 2 to request an amended report.

Alice K Tanner

Alice Tanner, Ph.D., M.S., CGC, FACMG, Laboratory Director

Genes and diseases tested

The personalized residual risks listed below are specific to this individual. The complete residual risk table is available at **go.sema4.com/residualrisk**

Table 1: List of genes and diseases tested with detailed results

	Disease	Gene	Inheritance Pattern	Status	Detailed Summary
⊖	**Negative**				
	Congenital Ichthyosis 4A and 4B	*ABCA12*	AR	Reduced Risk	**Personalized Residual Risk**: 1 in 1,500

AR=Autosomal recessive; XL=X-linked

Test methods and comments

Genomic DNA isolated from this patient was analyzed by one or more of the following methodologies, as applicable:

Fragile X CGG Repeat Analysis (Analytical Detection Rate >99%)

PCR amplification using Asuragen, Inc. AmplideX® *FMR1* PCR reagents followed by capillary electrophoresis for allele sizing was performed. Samples positive for *FMR1* premutations and full mutations greater than 90 CGG repeats in length were further analyzed by Southern blot analysis or methylation PCR to assess the size and methylation status of the *FMR1* CGG repeat. Additional testing to determine the status of AGG interruptions within the *FMR1* CGG repeat will be automatically performed for premutation alleles ranging from 55 to 90 repeats. These results, which may modify risk for expansion, will follow in a separate report.

Genotyping (Analytical Detection Rate >99%)

Multiplex PCR amplification and allele specific primer extension analyses using the MassARRAY® System were used to identify certain recurrent variants that are complex in nature or are present in low copy repeats. Rare sequence variants may interfere with assay performance.

Multiplex Ligation-Dependent Probe Amplification (MLPA) (Analytical Detection Rate >99%)

MLPA® probe sets and reagents from MRC-Holland were used for copy number analysis of specific targets versus known control samples. False positive or negative results may occur due to rare sequence variants in target regions detected by MLPA probes. Analytical sensitivity and specificity of the MLPA method are both 99%.

For alpha thalassemia, the copy numbers of the *HBA1* and *HBA2* genes were analyzed. Alpha-globin gene deletions, triplications, and the Constant Spring (CS) mutation are assessed. This test is expected to detect approximately 90% of all alpha-thalassemia mutations, varying by ethnicity. Carriers of alpha-thalassemia with three or more *HBA* copies on one chromosome, and one or no copies on the other chromosome, may not be detected. With the exception of triplications, other benign alpha-globin gene polymorphisms will not be reported. Analyses of *HBA1* and *HBA2* are performed in association with long-range PCR of the coding regions followed by short-read sequencing.

For Duchenne muscular dystrophy, the copy numbers of all *DMD* exons were analyzed. Potentially pathogenic single exon deletions and duplications are confirmed by a second method. Analysis of *DMD* is performed in association with sequencing of the coding regions.

For congenital adrenal hyperplasia, the copy number of the *CYP21A2* gene was analyzed. This analysis can detect large deletions typically due to unequal meiotic crossing-over between *CYP21A2* and the pseudogene *CYP21A1P*. Classic 30-kb deletions make up approximately 20% of *CYP21A2* pathogenic alleles. This test may also identify certain point mutations in *CYP21A2* caused by gene conversion events between *CYP21A2* and *CYP21A1P*. Some carriers may not be identified by dosage sensitive methods as this testing cannot detect individuals with two copies (duplication) of the *CYP21A2* gene on one chromosome and loss of *CYP21A2* (deletion) on the other chromosome. Analysis of *CYP21A2* is performed in association with long-range PCR of the coding regions followed by short-read sequencing.

For spinal muscular atrophy (SMA), the copy numbers of the *SMN1* and *SMN2* genes were analyzed. The individual dosage of exons 7 and 8 as well as the combined dosage of exons 1, 4, 6 and 8 of *SMN1* and *SMN2* were assessed. Copy number gains and losses can be detected with this assay. Depending on ethnicity, 6 - 29 % of carriers will not be identified by dosage sensitive methods as this testing cannot detect individuals with two copies (duplication) of the *SMN1* gene on one chromosome and loss of *SMN1* (deletion) on the other chromosome (silent 2+0 carrier) or individuals that carry an intragenic mutation in *SMN1*. Please also note that 2% of individuals diagnosed with SMA have a

causative *SMN1* variant that occurred de novo, and therefore cannot be picked up by carrier screening in the parents. Analysis of *SMN1* is performed in association with short-read sequencing of exons 2a-7, followed by confirmation using long-range PCR (described below).

In individuals with two copies of *SMN1* with Ashkenazi Jewish, East Asian, African American, Native American or Caucasian ancestry, the presence or absence of c.*3+80T>G significantly increases or decreases, respectively, the likelihood of being a silent 2+0 silent carrier.

MLPA for Gaucher disease *(GBA)*, cystic fibrosis *(CFTR)*, and non-syndromic hearing loss *(GJB2/GJB6)* will only be performed if indicated for confirmation of detected CNVs. If *GBA* analysis was performed, the copy numbers of exons 1, 3, 4, and 6 - 10 of the GBA gene (of 11 exons total) were analyzed. If *CFTR* analysis was performed, the copy numbers of all 27 *CFTR* exons were analyzed. If *GJB2/GJB6* analysis was performed, the copy number of the two *GJB2* exons were analyzed, as well as the presence or absence of the two upstream deletions of the *GJB2* regulatory region, del(*GJB6*-D13S1830) and del(*GJB6*-D13S1854).

Next Generation Sequencing (NGS) (Analytical Detection Rate >95%)

NGS was performed on a panel of genes for the purpose of identifying pathogenic or likely pathogenic variants.

Agilent SureSelect^{TM}XT Low Input technology was used with a custom capture library to target the exonic regions and intron/exon splice junctions of the relevant genes, as well as a number of UTR, intronic or promoter regions that contain previously reported mutations. Libraries were pooled and sequenced on the Illumina NovaSeq 6000 platform, using paired-end 100 bp reads. The sequencing data was analyzed using a custom bioinformatics algorithm designed and validated in house.

The coding exons and splice junctions of the known protein-coding RefSeq genes were assessed for the average depth of coverage (minimum of 20X) and data quality threshold values. Most exons not meeting a minimum of >20X read depth across the exon are further analyzed by Sanger sequencing. Please note that several genomic regions present difficulties in mapping or obtaining read depth >20X. These regions, which are described below, will not be reflexed to Sanger sequencing if the mapping quality or coverage is poor. Any variants identified during testing in these regions are confirmed by a second method and reported if determined to be pathogenic or likely pathogenic. However, as there is a possibility of false negative results within these regions, detection rates and residual risks for these genes have been calculated with the presumption that variants in these exons will not be detected, unless included in the MassARRAY® genotyping platform.

Exceptions: *ABCD1* (NM_000033.3) exons 8 and 9; *ACADSB* (NM_ 001609.3) chr10:124,810,695-124,810,707 (partial exon 9); *ADA* (NM_000022 2) exon 1; *ADAMTS2* (NM_014244 4) exon 1; *AGPS* (NM_003659.3) chr2:178,257,512-178,257,649 (partial exon 1); *ALDH7A1* (NM_001182.4) chr5:125,911,150-125,911,163 (partial exon 7) and chr5:125,896,807-125,896,821 (partial exon 10); *ALMS1* (NM_015120.4) chr2:73,612,990-73,613,041 (partial exon 1); *APOPT1* (NM_ 032374.4) chr14:104,040,437-104,040,455 (partial exon 3); *CDAN1* (NM_138477.2) exon 2; *CEP152* (NM_014985.3) chr15:49,061,146-49,061,165 (partial exon 14) and exon 22; *CEP290* (NM_025114.3) exon 5, exon 7, chr12:88,519,017-88,519,039 (partial exon 13), chr12:88,514,049-88,514,058 (partial exon 15), chr12:88,502,837-88,502,841 (partial exon 23), chr12:88,481,551-88,481,589 (partial exon 32), chr12:88,471,605-88,471,700 (partial exon 40); *CFTR* (NM_000492.3) exon 10; *COL4A4* (NM_000092 4) chr2:227,942,604-227,942,619 (partial exon 25); *COX10* (NM_001303.3) exon 6; *CYP11B1* (NM_000497.3) exons 3-7; *CYP11B2* (NM_000498.3) exons 3-7; *DNAI2* (NM_023036 4) chr17:72,308,136-72,308,147 (partial exon 12); *DOK7* (NM_173660.4) chr4:3,465,131-3,465,161 (partial exon 1) and exon 2; *DUOX2* (NM_014080.4) exons 6-8; *EIF2AK3* (NM_004836.5 exon 8; *EVC* (NM_153717 2) exon 1; *F5*(NM_000130 4) chr1:169,551,662-169,551,679 (partial exon 2); *FH* (NM_000143.3) exon 1; *GAMT* (NM_000156.5 exon 1; *GLDC*(NM_000170 2) exon 1; *GNPTAB* (NM_024312 4) chr17:4,837,000-4,837,400 (partial exon 2); *GNPTG* (NM_032520 4) exon 1; *GHR* (NM_000163 4) exon 3; *GYS2* (NM_021957.3) chr12:21,699,370-21,699,409 (partial exon 12); *HGSNAT* (NM_152419.2) exon 1; *IDS* (NM_000202.6 exon 3; *ITGB4* (NM_000213 4) chr17:73,749,976-73,750,060 (partial exon 33); *JAK3* (NM_000215.3) chr19:17,950,462-17,950,483 (partial exon 10); *LIFR* (NM_002310.5 exon 19; *LMBRD1* (NM_018368.3) chr6:70,459,226-70,459,257 (partial exon 5), chr6:70,447,828-70,447,836 (partial exon 7) and exon 12; *LYST* (NM_000081.3) chr1:235,944,158-235,944,176 (partial exon 16) and chr1 235,875,350-235,875,362 (partial exon 43); *MLYCD* (NM_012213.2) chr16:83,933,242-83,933,282 (partial exon 1); *MTR* (NM_000254.2) chr1 237,024,418-237,024,439 (partial exon 20) and chr1:237,038,019-237,038,029 (partial exon 24); *NBEAL2* (NM_015175 2) chr3 47,021,385-47,021,407 (partial exon 1); *NEB* (NM_001271208.1 exons 82-105; *NPC1* (NM_000271 4) chr18:21,123,519-21,123,538 (partial exon 14); *NPHP1* (NM_000272.3)chr2:110,937,251-110,937,263 (partial exon 3); *OCRL* (NM_000276.3) chrX:128,674,450-128,674,460 (partial exon 1); *PHKB* (NM_000293 2) exon 1 and chr16:47,732,498-47,732,504 (partial exon 30); *PIGN* (NM_176787.4) chr18:59,815,547-59,815,576 (partial exon 8); *PIP5K1C* (NM_012398.2) exon 1 and chr19:3637602-3637616 (partial exon 17); *POU1F1* (NM_000306.3) exon 5; *PTPRC* (NM_002838 4) exons 11 and 23; *PUS1* (NM_025215.5 chr12:132,414,446-132,414,532 (partial exon 2); *RPGRIP1L* (NM_015272 2) exon 23; *SGSH* (NM_000199.3) chr17:78,194,022-78,194,072 (partial exon 1); *SLC6A8* (NM_005629.3) exons 3 and 4; *ST3GAL5* (NM_003896.3) exon 1; *SURF1* (NM_003172.3) chrg:136,223,269-136,223,307 (partial exon 1); *TRPM6* (NM_017662 4) chrg:77,362,800-77,362,811 (partial exon 31); *TSEN54* (NM_207346.2) exon 1; *TYR* (NM_000372 4) exon 5; *VWF* (NM_000552.3) exons 24-26, chr12:6,125,675-6,125,684 (partial exon 30), chr12:6,121,244-6,121,265 (partial exon 33), and exon 34.

This test will detect variants within the exons and the intron-exon boundaries of the target regions. Variants outside these regions may not be detected, including, but not limited to, UTRs, promoters, and deep intronic areas, or regions that fall into the Exceptions mentioned above. This technology may not detect all small insertion/deletions and is not diagnostic for repeat expansions and structural genomic variation. In addition, a mutation(s) in a gene not included on the panel could be present in this patient.

Variant interpretation and classification was performed based on the American College of Medical Genetics Standards and Guidelines for the Interpretation of Sequence Variants (Richards et al, 2015). All potentially pathogenic variants may be confirmed by either a specific genotyping assay or Sanger sequencing, if indicated. Any benign variants, likely benign variants or variants of uncertain significance identified during this analysis will not be reported.

Next Generation Sequencing for *SMN1*

Exonic regions and intron/exon splice junctions of *SMN1* and *SMN2* were captured, sequenced, and analyzed as described above. Any variants located within exons 2a-7 and classified as pathogenic or likely pathogenic were confirmed to be in either *SMN1* or *SMN2* using gene-specific long-range PCR analysis followed by Sanger sequencing. Variants located in exon 1 cannot be accurately assigned to either *SMN1* or *SMN2* using our current methodology, and so these variants are not reported.

Copy Number Variant Analysis (Analytical Detection Rate >95%)

Large duplications and deletions were called from the relative read depths on an exon-by-exon basis using a custom exome hidden Markov model (XHMM) algorithm. Deletions or duplications determined to be pathogenic or likely pathogenic were confirmed by either a custom arrayCGH platform, quantitative PCR, or MLPA (depending on CNV size and gene content). While this algorithm is designed to pick up deletions and duplications of 2 or more exons in length, potentially pathogenic single-exon CNVs will be confirmed and reported, if detected. Deletions and duplications near the lower limit of detection may not be detected due to run variability.

Exon Array (Confirmation method) (Accuracy >99%)

The customized oligonucleotide microarray (Oxford Gene Technology) is a highly-targeted exon-focused array capable of detecting medically relevant microdeletions and microduplications at a much higher resolution than traditional aCGH methods. Each array matrix has approximately 180,000 60-mer oligonucleotide probes that cover the entire genome. This platform is designed based on human genome NCBI Build 37 (hg19) and the CGH probes are enriched to target the exonic regions of the genes in this panel.

Quantitative PCR (Confirmation method) (Accuracy >99%)

The relative quantification PCR is utilized on a Roche Universal Library Probe (UPL) system, which relates the PCR signal of the target region in one group to another. To test for genomic imbalances, both sample DNA and reference DNA is amplified with primer/probe sets that specific to the target region and a control region with known genomic copy number. Relative genomic copy numbers are calculated based on the standard ΔΔCt formula.

Long-Range PCR (Analytical Detection Rate >99%)

Long-range PCR was performed to generate locus-specific amplicons for *CYP21A2*, *HBA1* and *HBA2* and *GBA*. The PCR products were then prepared for short-read NGS sequencing and sequenced. Sequenced reads were mapped back to the original genomic locus and run through the bioinformatics pipeline. If indicated, copy number from MLPA was correlated with the sequencing output to analyze the results. Please note that in rare cases, allele drop-out may occur, which has the potential to lead to false negative results. For *CYP21A2*, a certain percentage of healthy individuals carry a duplication of the *CYP21A2* gene, which has no clinical consequences. In cases where multiple copies of *CYP21A2* are located on the same chromosome in tandem, only the last copy will be amplified and assessed for potentially pathogenic variants, due to size limitations of the PCR reaction. However, because these alleles contain at least two copies of the *CYP21A2* gene in tandem, it is expected that this patient has at least one functional gene in the tandem allele and this patient is therefore less likely to be a carrier. A *CYP21A1P/ CYP21A2* hybrid gene detected only by MLPA but not by long-range PCR will not be reported when the long-range PCR indicates the presence of two full *CYP21A2* gene copies (one on each chromosome), as the additional hybrid gene is nonfunctional. Classic 30-kb deletions are identified by MLPA and are also identified by the presence of multiple common pathogenic *CYP21A2* variants by long-range PCR. Since multiple pseudogene-derived variants are detected in all cases with the classic 30kb deletion, we cannot rule out the possibility that some variant(s) detected could be present in trans with the chimeric *CYP21A1P/ CYP21A2* gene created by the 30kb deletion. When an individual carries both a duplication allele and a pathogenic variant, or multiple pathogenic variants, the current analysis may not be able to determine the phase (cis/trans configuration) of the *CYP21A2* alleles identified. Family studies may be required in certain scenarios where phasing is required to determine the carrier status.

Residual Risk Calculations

Carrier frequencies and detection rates for each ethnicity were calculated through the combination of internal curations of >30,000 variants and genomic frequency data from >138,000 individuals across seven ethnic groups in the gnomAD database. Additional variants in HGMD and novel deleterious variants were also incorporated into the calculation. Residual risk values are calculated using a Bayesian analysis combining the a *priori* risk of being a pathogenic mutation carrier (carrier frequency) and the detection rate. They are provided only as a guide for assessing approximate risk given a negative result, and values will vary based on the exact ethnic background of an individual. This report does not represent medical advice but should be interpreted by a genetic counselor, medical geneticist or physician skilled in genetic result interpretation and the relevant medical literature.

Personalized Residual Risk Calculations

Agilent SureSelect^{TM}XT Low-Input technology was utilized in order to create whole-genome libraries for each patient sample. Libraries were then pooled and sequenced on the Illumina NovaSeq platform. Each sequencing lane was multiplexed to achieve 0.4-2x genome coverage, using paired-end 100 bp reads. The sequencing data underwent ancestral analysis using a customized, licensed bioinformatics algorithm that was validated in house. Identified sub-ethnic groupings were binned into one of 7 continental-level groups (African, East Asian, South Asian, Non-Finnish European, Finnish, Native American, and Ashkenazi Jewish) or, for those ethnicities that matched poorly to the continental-level groups, an 8th "unassigned" group, which were then used to select residual risk values for each gene. For individuals belonging to multiple high-level ethnic groupings, a weighting strategy was used to select the most appropriate residual risk. For genes that had insufficient data to calculate ethnic-specific residual risk values, or for sub-ethnic groupings that fell into the "unassigned" group, a "worldwide" residual risk was used. This "worldwide" residual risk was calculated using data from all available continental-level groups.

Several genes have multiple residual risks associated to reflect the likelihood of the tested individual being a carrier for different diseases that are attributed to non-overlapping pathogenic variants in that gene. When calculating the couples' combined reproductive risk, the highest residual risk for each patient was selected.

Sanger Sequencing (Confirmation method) (Accuracy >99%)

Sanger sequencing, as indicated, was performed using BigDye Terminator chemistry with the ABI 3730 DNA analyzer with target specific amplicons. It also may be used to supplement specific guaranteed target regions that fail NGS sequencing due to poor quality or low depth of coverage (<20 reads) or as a confirmatory method for NGS positive results. False negative results may occur if rare variants interfere with amplification or annealing.

Tay-Sachs Disease (TSD) Enzyme Analysis (Analytical Detection Rate ≥98%)

Hexosaminidase activity and Hex A% activity were measured by a standard heat-inactivation, fluorometric method using artificial 4-MU-β-N-acetyl glucosaminide (4-MUG) substrate. This assay is highly sensitive and accurate in detecting Tay-Sachs carriers and individuals affected with TSD. Normal ranges of Hex A% activity are 55.0-72.0 for white blood cells and 58.0-72.0 for plasma. It is estimated that less than 0.5% of Tay-Sachs carriers have non-carrier levels of percent Hex A activity, and therefore may not be identified by this assay. In addition, this assay may detect individuals that are carriers of or are affected with Sandhoff disease. False positive results may occur if benign variants, such as pseudodeficiency alleles, interfere with the enzymatic assay. False negative results may occur if both *HEXA* and *HEXB* pathogenic or pseudodeficiency variants are present in the same individual.

Please note that it is not possible to perform Tay-Sachs disease enzyme analysis on saliva samples, buccal swabs, tissue samples, semen samples, or on samples received as extracted DNA.

This test was developed, and its performance characteristics determined by Sema4 Opco, Inc. It has not been cleared or approved by the US Food and Drug Administration. FDA does not require this test to go through premarket FDA review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments (CLIA) as qualified to perform high complexity clinical laboratory testing. These analyses generally provide highly accurate information regarding the patient's carrier or affected status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.

SELECTED REFERENCES

Carrier Screening

Grody W et al. ACMG position statement on prenatal/preconception expanded carrier screening. *Genet Med.* 2013 15:482-3.

Fragile X syndrome:

Chen L et al. An information-rich CGG repeat primed PCR that detects the full range of Fragile X expanded alleles and minimizes the need for Southern blot analysis. *J Mol Diag* 2010 12:589-600.

Spinal Muscular Atrophy:

Luo M et al. An Ashkenazi Jewish *SMN1* haplotype specific to duplication alleles improves pan-ethnic carrier screening for spinal muscular atrophy. *Genet Med.* 2014 16:149-56.

Ashkenazi Jewish Disorders:

Scott SA et al. Experience with carrier screening and prenatal diagnosis for sixteen Ashkenazi Jewish Genetic Diseases. *Hum. Mutat.* 2010 31:1-11.

Akler G et al. Towards a unified approach for comprehensive reproductive carrier screening in the Ashkenazi, Sephardi, and Mizrahi Jewish populations. *Mol Genet Genomic Med.* 2020 Feb 8(2):e1053.

Duchenne Muscular Dystrophy:

Flanigan KM et al. Mutational spectrum of *DMD* mutations in dystrophinopathy patients: application of modern diagnostic techniques to a large cohort. *Hum Mutat.* 2009 30:1657-66.

Variant Classification:

Richards S et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med.* 2015 May:17(5):405-24

Additional disease-specific references available upon request.

LabCorp
Laboratory Corporation of America

Client/Sending Facility:
Seattle Sperm Bank

4915 25th Ave Ne Ste 204
SEATTLE, WA 98105
Ph: (206)588-1484
Fax: (206) 466-4696 WAB-55

LCLS Specimen Number:	**056-129-1154-0**	Account Number:	46857540
Patient Name:	**12552, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	02/25/2020
Lab Number:	DCGWA20-701 L	Date Received:	02/26/2020
Indications:	DONOR	Date Reported:	**03/09/2020**

Test: **Chromosome, Blood, Routine**

Cells Counted:	15	Cells Karyotyped:	2
Cells Analyzed:	5	Band Resolution:	550

CYTOGENETIC RESULT: 46,XY

INTERPRETATION: NORMAL MALE KARYOTYPE

Cytogenetic analysis of PHA stimulated cultures has revealed a MALE karyotype with an apparently normal GTW banding pattern in all cells observed.

This result does not exclude the possibility of subtle rearrangements below the resolution of cytogenetics or congenital anomalies due to other etiologies.

LabCorp
Laboratory Corporation of America

Client/Sending Facility:
Seattle Sperm Bank

4915 25th Ave Ne Ste 204
SEATTLE, WA 98105
Ph: (206)588-1484
Fax: (206) 466-4696 WAB-55

LCLS Specimen Number:	**056-129-1154-0**
Patient Name:	**12552, DONOR**
Date of Birth:	[redacted]
Gender:	M
Patient ID:	
Lab Number:	DCGWA20-701 L

Account Number:	46857540
Ordering Physician:	**J OLLIFFE**
Specimen Type:	**BLOOD**
Client Reference:	
Date Collected:	02/25/2020
Date Received:	02/26/2020



LabCorp
Laboratory Corporation of America

Client/Sending Facility:
Seattle Sperm Bank

4915 25th Ave Ne Ste 204
SEATTLE, WA 98105
Ph: (206)588-1484
Fax: (206) 466-4696 WAB-55

LCLS Specimen Number:	**056-129-1154-0**
Patient Name:	**12552, DONOR**
Date of Birth:	[redacted]
Gender:	M
Patient ID:	
Lab Number:	DCGWA20-701 L

Account Number:	46857540
Ordering Physician:	**J OLLIFFE**
Specimen Type:	**BLOOD**
Client Reference:	
Date Collected:	02/25/2020
Date Received:	02/26/2020

Hiba Risheg, PhD., FACMG

Daniel Toweill, MD
Medical Director
Stuart Schwartz, PhD
National Director of Cytogenetics

Technical component performed by Laboratory Corporation of America Holdings, 550 17th Ave Ste 300 , Seattle , WA , 98122-5789 (206) 861-7000
Professional Component performed by LabCorp CLIA 50D0630157, 550 17th Ave. Suite 200, Seattle WA 98122-5789. Medical Director, Daniel Toweill, MD
This document contains private and confidential health information **protected by state and federal law.**

LabCorp

Patient Report

Specimen ID: 056-129-2118-0
Control ID: B0099603720

Acct #: 46857540 **Phone:** (206) 588-1484 **Rte:** 00

12552, DONOR

Seattle Sperm Bank
4915 25th Ave Ne Ste 204
SEATTLE WA 98105

Patient Details
DOB: [redacted]
Age(y/m/d): [redacted]
Gender: M **SSN:**
Patient ID:

Specimen Details
Date collected: 02/25/2020 0740 Local
Date received: 02/26/2020
Date entered: 02/25/2020
Date reported: 03/02/2020 1506 ET

Physician Details
Ordering: J OLLIFFE
Referring:
ID:
NPI: 1306838271

General Comments & Additional Information
Alternate Control Number: B0099603720

Alternate Patient ID: Not Provided

Ordered Items
LP+12AC+CBC/D/Plt+UA+Rh+ABO...

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
LP+12AC+CBC/D/Plt+UA+Rh+ABO...					
Chemistries					01
Glucose	93		mg/dL	65 - 99	01
Uric Acid	6.0		mg/dL	3.7 - 8.6	01
Please Note:					01
	Therapeutic target for gout patients: <6.0				
BUN	11		mg/dL	6 - 20	01
Creatinine	0.98		mg/dL	0.76 - 1.27	01
eGFR If NonAfricn Am	102		mL/min/1.73	>59	
eGFR If Africn Am	118		mL/min/1.73	>59	
Calcium	9.3		mg/dL	8.7 - 10.2	01
Protein, Total	7.2		g/dL	6.0 - 8.5	01
Albumin	4.9		g/dL	4.0 - 5.0	01
	Please note reference interval change				
Bilirubin, Total	0.4		mg/dL	0.0 - 1.2	01
Alkaline Phosphatase	68		IU/L	39 - 117	01
LDH	157		IU/L	121 - 224	01
AST (SGOT)	15		IU/L	0 - 40	01
ALT (SGPT)	17		IU/L	0 - 44	01
					01
Lipids					01
Cholesterol, Total	198		mg/dL	100 - 199	01
Triglycerides	139		mg/dL	0 - 149	01
HDL Cholesterol	50		mg/dL	>39	01
LDL Cholesterol Calc	**120**	**High**	mg/dL	0 - 99	
LDL/HDL Ratio	2.4		ratio	0.0 - 3.6	
Please Note:					01

LDL/HDL Ratio

	Men	Women
1/2 Avg.Risk	1.0	1.5
Avg.Risk	3.6	3.2
2X Avg.Risk	6.2	5.0

13 MAR 20

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error, please call 800-598-3345

© 1995-2020 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version: 1.00

LabCorp **Patient Report**

Patient: 12552, DONOR
DOB: [redacted] **Patient ID:**
Control ID: B0099603720
Specimen ID: 056-129-2118-0
Date collected: 02/25/2020 0740 Local

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
			3X Avg.Risk	8.0 6.1	
					01
Hemoglobin Fractionation					01
Hgb F	0.0		%	0.0 - 2.0	02
Hgb A	98.0		%	96.4 - 98.8	02
Hgb S	0.0		%	0.0	02
Hgb C	0.0		%	0.0	02
Hgb A2	2.0		%	1.8 - 3.2	02
Hgb Variant	0.0		%	0.0	02
Interpretation					
Normal adult hemoglobin present.					02
ABO Grouping	A				01
Rh Factor	Positive				01
Please note: Prior records for this patient's ABO / Rh type are not available for additional verification.					
					01
CBC, Platelet Ct, and Diff					01
WBC	5.1		x10E3/uL	3.4 - 10.8	01
RBC	5.57		x10E6/uL	4.14 - 5.80	01
Hemoglobin	14.4		g/dL	13.0 - 17.7	01
Hematocrit	45.2		%	37.5 - 51.0	01
MCV	81		fL	79 - 97	01
MCH	**25.9**	Low	pg	26.6 - 33.0	01
MCHC	31.9		g/dL	31.5 - 35.7	01
RDW	13.6		%	11.6 - 15.4	01
Platelets	302		x10E3/uL	150 - 450	01
Neutrophils	64		%	Not Estab.	01
Lymphs	28		%	Not Estab.	01
Monocytes	7		%	Not Estab.	01
Eos	1		%	Not Estab.	01
Basos	0		%	Not Estab.	01
Neutrophils (Absolute)	3.2		x10E3/uL	1.4 - 7.0	01
Lymphs (Absolute)	1.4		x10E3/uL	0.7 - 3.1	01
Monocytes(Absolute)	0.4		x10E3/uL	0.1 - 0.9	01
Eos (Absolute)	0.1		x10E3/uL	0.0 - 0.4	01
Baso (Absolute)	0.0		x10E3/uL	0.0 - 0.2	01
Immature Granulocytes	0		%	Not Estab.	01
Immature Grans (Abs)	0.0		x10E3/uL	0.0 - 0.1	01
					01
Urinalysis Gross Exam					01
Specific Gravity	1.030			1.005 - 1.030	01
pH	6.0			5.0 - 7.5	01
Urine-Color	Yellow			Yellow	01

13 Mar 20

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error, please call 800-598-3345

© 1995-2020 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version: 1.00